

Treatment for comorbid depressive disorder or subthreshold depression in diabetes mellitus: Systematic review and meta-analysis

Christina van der Feltz-Cornelis¹  | Sarah F. Allen¹  | Richard I. G. Holt²  |
Richard Roberts³ | Arie Nouwen⁴  | Norman Sartorius⁵ 

¹Department of Health Sciences, Hull York Medical School, University of York, York, UK

²Human Development and Health, Faculty of Medicine, University of Southampton, Southampton, UK

³Department of Family Medicine & Community Health, University of Wisconsin, Madison, WI, USA

⁴Department of Psychology, Middlesex University, London, UK

⁵Association for the Improvement of Mental Health Programmes, Geneva, Switzerland

Correspondence

Christina van der Feltz-Cornelis, Mental Health and Addictions Research Group, Department of Health Sciences, HYMS, University of York, Heslington, York, YO105DD, UK.

Email: Christina.vanderfeltz-Cornelis@york.ac.uk

Funding information

Hull York Medical School, Grant/Award Number: NA

Abstract

Objective: To provide an estimate of the effect of interventions on comorbid depressive disorder (MDD) or subthreshold depression in type 1 and type 2 diabetes.

Methods: Systematic review and meta-analysis. We searched PubMed, PsycINFO, Embase, and the Cochrane Library for randomized controlled trials evaluating the outcome of depression treatments in diabetes and comorbid MDD or subthreshold symptoms published before August 2019 compared to care as usual (CAU), placebo, waiting list (WL), or active comparator treatment as in a comparative effectiveness trial (CET). Primary outcomes were depressive symptom severity and glycemic control. Cohen's d is reported.

Results: Forty-three randomized controlled trials (RCTs) were selected, and 32 RCTs comprising 3,543 patients were included in the meta-analysis. Our meta-analysis showed that, compared to CAU, placebo or WL, all interventions showed a significant effect on combined outcome 0.485 (95% CI 0.360; 0.609). All interventions showed a significant effect on depression. Pharmacological treatment, group therapy, psychotherapy, and collaborative care had a significant effect on glycemic control. High baseline depression score was associated with a greater reduction in HbA_{1c} and depressive outcome. High baseline HbA_{1c} was associated with a greater reduction in HbA_{1c}.

Conclusion: All treatments are effective for comorbid depression in type 1 diabetes and type 2 diabetes. Over the last decade, new interventions with large effect sizes have been introduced, such as group-based therapy, online treatment, and exercise. Although all interventions were effective for depression, not all treatments were effective for glycemic control. Effective interventions in comorbid depressive disorder may not be as effective in comorbid subthreshold depression. Baseline depression and HbA_{1c} scores modify the treatment effect. Based on the findings, we provide

Registration: International Prospective Register of Systematic Reviews (PROSPERO) number CRD42019147910

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Brain and Behavior* published by Wiley Periodicals LLC

guidance for treatment depending on patient profile and desired outcome, and discuss possible avenues for further research.

KEYWORDS

depression, diabetes mellitus, glycemic control, meta-analysis, systematic review, treatments

1 | SUMMATIONS

This systematic review and meta-analysis exploring psychotherapeutic, pharmacologic, and other interventions shows beneficial treatment effects for comorbid depression in type 1 and type 2 diabetes mellitus with moderate-to-large effect sizes for most intervention types.

Although all interventions were effective for depression, not all treatments were effective for glycemic control.

Effective interventions in comorbid depressive disorder may not be as effective in comorbid subthreshold depression.

2 | LIMITATIONS

Most of the selected studies did not meet all criteria to reduce the risk of bias and not all provided sufficient data to be included in the meta-analysis.

Further, some treatments were only evaluated in a single RCT.

There is a scarcity of data from many low- and middle-income countries.

3 | INTRODUCTION

No international consensus exists to guide treatment of comorbid depression in diabetes. Nonetheless, over the last three decades, clinicians have been seeing increasing numbers of patients with comorbid depression of various severity in diabetes (Khaledi et al., 2019; Zheng et al., 2018) due to the exploding prevalence of both diabetes and depression (GBD Disease & Injury Incidence & Prevalence Collaborators, 2018). This can amount to up to 30% depending on severity of symptoms and it occurs especially where the person with diabetes has elevated HbA_{1c} despite treatment, or frequent episodes of hypoglycemia and increased glucose variability, diabetes-related complications, and disengagement from treatments (Groot et al., 2001; Lustman, Anderson, et al., 2000; O'Connor et al., 2009). Depression is a serious psychiatric disorder characterized by loss of interest or pleasure, depressed mood, and suicidal behavior (Ruengorn et al., 2012). Diabetes and depression can both seriously affect an individual's quality of life, and lead to functional disability, increased distress, and social burden (Renn et al., 2011). Depressive symptoms in people with diabetes can have a detrimental impact on engagement with diabetes management (Ciechanowski et al., 2000; Gonzalez, Peyrot, et al., 2008) and on glycemic control (Lustman, Anderson, et al., 2000) as well as on health-related outcomes (e.g.,

weight gain and diabetes-related complications) and associated healthcare costs (Black et al., 2003). As such, the high prevalence of this comorbidity is accompanied by high rates of morbidity and mortality worldwide (Hofmann et al., 2013; Lloyd et al., 2018; Nouwen et al., 2019). Epidemiological studies indicate there is a bidirectional relationship between diabetes and depression (Golden et al., 2008; Katon, 2008; Katon et al., 2007), in which individuals with diabetes have an increased risk of depression and vice versa; the presence of a depressive disorder can increase the risk of metabolic diseases such as diabetes (Renn et al., 2011) and there is an association between depression and diabetes complications (Groot et al., 2001; Van Steenberg-Weijnenburg et al., 2011).

Evidence is growing to suggest that depression may play a role in the pathogenesis of diabetes in a number of ways. Depression may be a consequence of similar environmental factors that govern glucose metabolism, and can also independently influence nutrition and lifestyle choices which can predispose individuals to the development of diabetes (Beydoun & Wang, 2010). Biological mechanisms have also been proposed through a dysregulated and overactive HPA axis, a shift in sympathetic nervous system tone toward enhanced sympathetic activity, and a pro-inflammatory state (Champaneri et al., 2010; Joseph & Golden, 2017). The role of inflammation is particularly pertinent. Laake et al. (2014) found that increased inflammation may be involved in the pathogenesis of depression in people with type 2 diabetes, which in turn could contribute to the increased risk of complications and mortality in this clinical population (Geraets et al., 2020).

The relationship between depressive symptoms and poorer diabetes self-care (Gonzalez, Safren, et al., 2008) applies also to subclinical or subthreshold depressive symptoms (Pibernik-Okanović et al., 2011) and not only to major depressive disorder. Subthreshold refers to those with two or more depressive symptoms who do not meet the diagnostic criteria for depression (Rodríguez et al., 2012). Subthreshold depressive symptoms in people with diabetes have been found to be persistent but also associated with an increased risk of worsening over time (Bot et al., 2010; Nefs et al., 2012; Pibernik-Okanovic et al., 2008). Furthermore, an increased incidence of adverse health outcomes and suboptimal metabolic control has been observed not only in patients with the established diagnosis of depression but also in those suffering subthreshold depressive symptoms (Johnson et al., 2014). This indicates that even mild depression is clinically relevant, and implies that combined treatments could also be efficacious for people with diabetes and subthreshold depressive symptoms.

A lack of a clear understanding of the shared origins of depression and diabetes means that finding the most appropriate treatment

for this comorbidity in this vulnerable patient group is difficult. In order to optimize health outcomes, feasible and effective interventions aiming to provide benefits to both physical and mental health are recommended (Baumeister & Bengel, 2012; Baumeister et al., 2014; Harkness et al., 2010). The focus of treatment strategies should be on the remission or improvement of depression, in addition to improvement in glycemic control as a marker of diabetes outcome (Petrak et al., 2015).

Evidence shows that comorbid depression in diabetes can be treated with moderate success by psychological and pharmacological interventions, often implemented by using collaborative care (Katon, Von Korff, et al., 2004) and stepped care approaches (Stoop et al., 2015). However, there is conflicting evidence for the efficacy of antidepressants and psychological therapy in the improvement of glycemic control (Lustman, Anderson, et al., 2000; Lustman et al., 1997, 1998a, 2000b, 2007). Petrak, Herpertz, et al. (2015) claim that more research is needed to evaluate treatment of different subtypes of depression in people with diabetes as well as the effectiveness of new approaches to treatment.

3.1 | Rationale and objective

A previous systematic review of treatments for comorbid depression in diabetes indicated favorable effects on depressive outcome according to rating scales (Van der Feltz-Cornelis et al., 2010), but did not include data for subthreshold depression, which has been found to be related to poorer diabetes outcomes similar to DSM-5 depressive disorder (Gonzalez, Safren, et al., 2008; Pibernik-Okanović et al., 2011). We updated and expanded this systematic review and meta-analysis of randomized controlled trials to provide an estimate of the effect of interventions for comorbid depressive disorder or subthreshold depression in type 1 diabetes and type 2 diabetes. The interventions were compared with care as usual (CAU), waiting list (WL), placebo or another active comparator (e.g., another antidepressant or psychotherapy) on depression outcome and glycemic control, and, if possible, to provide treatment guidance for this condition.

4 | METHOD

This systematic review and meta-analysis was performed in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement (Liberati et al., 2009). We searched MEDLINE, Embase, the Cochrane Central Register of Controlled Trials, and Web of Science using Ovid software. The full search strategy and keywords used have been published elsewhere (Van der Feltz-Cornelis et al., 2010) and are shown in the appendix (pp 1-2). The reference lists of selected RCTs and reviews were checked for relevant studies that were not included in the databases. The search was supported by the Centre for Reviews and Dissemination at the University of York.

The protocol for this review is registered on PROSPERO and can be found here: https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42019147910

The final search results were restricted to studies completed before 28th August 2019. Inclusion criteria for studies were that they should be randomized clinical trials, provide a treatment intended to have an effect on both comorbid depressive symptoms and glycemic control in type 1 diabetes and/or type 2 diabetes, and have a control arm (e.g., CAU, placebo, WL or active comparator). The intervention had to be described sufficiently in order to be classified as a psychotherapeutic, medical, pharmacological, collaborative care or other type of intervention. A glossary providing an explanation about the interventions and a list of acronyms are provided in the appendix (pp 3-4).

Participants were adult patients with diabetes and comorbid depressive or subthreshold depression, which was defined as the presence of two or more core depressive symptoms, but not meeting the DSM-5 diagnostic criteria for depressive disorder (Rodríguez et al., 2012). No restriction was placed on type of intervention or publication language. Studies were not included if depressive disorder or depressive symptoms were not established in a systematic manner such as by semistructured interview or questionnaire at baseline. Studies were selected in a two-stage process. First, titles and abstracts from the electronic searches were scrutinized by two independent reviewers (SA and CFC). Second, if the abstract met inclusion criteria, we obtained full texts and final decisions were made about study inclusion. Disagreement regarding inclusion status was discussed. Consensus was reached in all cases.

Two reviewers (SA and CFC) independently extracted data for participants' characteristics, interventions, and study outcomes. A proforma as used in the original systematic review (Van der Feltz-Cornelis et al., 2010) was used to extract data from the included studies, now also including subthreshold depression from the search hits. The extracted data included: author and year; country; study type; sample size; age; baseline depression measure/diagnostic tool; baseline depression score, baseline glycemic control score, intervention details; control group, length of follow-up; diabetes and depression outcomes with regard to: i) the change in depression score from baseline to last follow-up using any validated self-report measure of depressive symptomatology and ii) the change in levels of biological marker of glycemic control from baseline to last follow-up. Assessment of glycemic control could be using HbA_{1c}, which provides an integrated measure of mean blood glucose levels over the last 6-8 weeks, or FBG, which gives an indication of the blood glucose concentration at the moment of assessment. If both were reported, we used the HbA_{1c} to calculate a standardized mean difference. The difference in means of each outcome was the primary measure within each study. Additional outcomes on adherence to recommendations of health-care providers with regard to self-care behaviors were extracted if reported. Authors were approached for additional data when questions arose.

4.1 | Risk-of-bias assessment

The Cochrane risk-of-bias tool (McGuire et al., 1998) was used to assess random sequence generation (selection bias), allocation concealment (selection bias), blinding of participants and personnel (performance bias), blinding of outcome assessment (detection bias), incomplete outcome data (attrition bias), selective reporting (reporting bias), and other biases. Risk of bias was assessed by SA and CFC independently.

Initial disagreements were resolved by consensus (Appendix pp 6–9). As psychotherapy trials often have limitations in the possibility for blinding (Van der Feltz-Cornelis & Ader, 2000), studies with limited blinding procedures were not excluded from the analysis. GRADE assessments were made (Guyatt et al., 2008) to give the confidence in each reported effect size. They are reported in the appendix (Appendix page 6–9).

4.2 | Statistical analysis

As a first step, overall meta-analysis was performed for all RCTs comparing all treatments with CAU, WL, or placebo for the combined effect on depressive outcome and glycemic control (illness burden). Then, we performed an analysis of illness burden in the studies reporting on depression versus the studies reporting on subthreshold depression. Then, studies were grouped according to the mode of treatment (pharmacotherapy, psychotherapy, collaborative care, online, phone and group interventions, exercise), depression severity (both as depression scores at baseline, and as classification of major depressive disorder or subthreshold depression), and depressive or diabetes outcome. Effect sizes were calculated. Outcomes from individual studies were pooled using a random-effects model (DerSimonian & Laird, 1986), as this approach assumes that there could be clinical and methodological heterogeneity that might affect the findings. All pooled analyses were reported with 95% confidence intervals (CIs). The effects were presented in terms of standardized effect sizes (Cohen's *d*). An effect size of 0.5 indicates that the mean of the experimental group is half a standard unit larger than the mean of the control group. It is generally assumed that an effect size of 0.56–1.2 represents a large clinical effect, while effect sizes of 0.33–0.55 are moderate, and effect sizes of 0–0.32 are small (Lipsey & Wilson, 1993). A meta-regression was conducted to assess whether baseline levels of depressive severity (scores on depression questionnaires) (Appendix pp.15) or glycemic control (HbA_{1c}) influenced the effect of the intervention. Between-study heterogeneity was assessed using the *I* (Khalidi et al., 2019) statistic (Higgins, 2003). Publication bias was examined by constructing a Begg funnel plot (Begg, 1994) and Duval's trim and fill (Rothstein et al., 2005). We adhered to published guidance of the Cochrane handbook (Higgins et al., 2019) throughout. We used the statistical program Comprehensive Meta-Analysis, version 2 (Biostat, 2005) to conduct random-effects meta-analyses.

5 | RESULTS

A PRISMA flowchart of study selection is presented in the appendix (pp 5). The overall search strategy yielded 8,684 citations of which 43 studies with 4,602 patients were included. This included fifteen studies from the original systematic review (Echeverry et al., 2009; Ell et al., 2010; Gulseren et al., 2005; Huang et al., 2002; Katon, Von Korff, et al., 2004; Li et al., 2003; Lu Xs & Bx, 2005; Lustman, Freedland, et al., 2000; Lustman et al., 1997, 1998b; Paile-Hyvärinen et al., 2003; Paile-Hyvärinen et al., 2007; Simson et al., 2008; Williams et al., 2004; Xue, 2004). Of the selected 43 studies, 39 were written or available in English, and four in Chinese (Huang et al., 2002; Li et al., 2003; Lu Xs & Bx, 2005; Xue, 2004). The latter were translated by certified translators and were included in the review. Eight trials with active comparator (CETs) were not entered in the meta-analysis in accordance with the Cochrane Handbook instructions for dealing with heterogeneity (Higgins et al., 2019), as pooling was not possible because of heterogeneous control groups (Barragán-Rodríguez et al., 2008; Gois et al., 2014; Gulseren et al., 2005; Kang et al., 2015; Karaiskos et al., 2013; Khazaie et al., 2011; Kumar et al., 2015; Petrak, Herpertz, et al., 2015). Three RCTs were not entered because they did not present the data required for pooling (Bastelaar et al., 2011; Brouwer et al., 2019; Ell et al., 2011). Thirty-two RCTs with 3,543 patients with type 1 diabetes and type 2 diabetes were entered into the meta-analysis (Bogner et al., 2012; Ebert et al., 2017; Echeverry et al., 2009; Ell et al., 2010; Groot et al., 2019; Guo et al., 2014; Hermanns et al., 2015; Huang et al., 2002, 2016; Johnson et al., 2014; Katon, Von Korff, et al., 2004; Li et al., 2003; Long et al., 2015; Lu Xs & Bx, 2005; Lustman, Freedland, et al., 2000; Lustman et al., 1997, 1998b; Naik et al., 2019; Newby et al., 2017; Paile-Hyvärinen et al., 2003; Paile-Hyvärinen et al., 2007; Penckofer et al., 2012; Pibernik-Okanovic et al., 2009, 2015; Piette et al., 2011; Safren et al., 2014; Schneider et al., 2016; Simson et al., 2008; Tovote et al., 2014; Williams et al., 2004; Xue, 2004; Zheng et al., 2015) All studies were performed in patients with diabetes as the primary or index condition, who suffered from comorbid depressive disorder or subthreshold depression. The countries in which each study was conducted are shown in Figure 1.

Results are shown in Table 1. The studies reported mostly on type 2 diabetes, or on type 1 and type 2 diabetes combined.

Overall meta-analysis in the RCTs comparing all treatments with CAU, WL, or placebo for the combined effect on depressive outcome and glycemic control showed an effect size of 0.485; 95% CI 0.360; 0.609, $p < .0001$ (Appendix pp 10–12).

Twenty-four studies (Atlantis et al., 2014; Bogner et al., 2012; Echeverry et al., 2009; Groot et al., 2019; Guo et al., 2014; Huang et al., 2002; Johnson et al., 2014; Katon, Von Korff, et al., 2004; Li et al., 2003; Long et al., 2015; Lu Xs & Bx, 2005; Lustman, Freedland, et al., 2000; Lustman et al., 1998b; Naik et al., 2019; Newby et al., 2017; Paile-Hyvärinen et al., 2003; Paile-Hyvärinen et al., 2007; Piette et al., 2011; Safren et al., 2014; Schneider et al., 2016; Simson et al., 2008; Williams et al., 2004; Xue, 2004) examined patients with diabetes and depressive disorder, termed major depressive disorder




FIGURE 1 Map of the world showing the location of each study included in the review

(MDD). All treatments showed significant effects in terms of depression outcomes. Large effect sizes were found in group-based therapy: effect size 1.650 (95% CI 1.196; 2.103), $p = .0001$; online treatment: effect size 0.789 (95% CI 0.358; 1.219), $p = .0001$; exercise: effect size 0.648 (95% CI 0.120; 1.177), $p = .016$; pharmacological treatment: effect size 0.571 (95% CI 0.348; 0.794) $p = .0001$, and psychotherapy: effect size 0.558 (95% CI 0.417; 0.700), $p = .0001$. Moderate effect sizes were found in collaborative care: effect size 0.434 (95% CI 0.284; 0.583), $p < .0001$; and phone treatment: effect size 0.344 (95% CI 0.034; 0.654), $p = .030$. The forest plot is shown in Figure 2 below.

Treatment showed a significant but small effect size in terms of glycemic control: 0.208 (95% CI 0.088; 0.329), $p = .001$. However, the effect size differed between treatment types: pharmacological treatment 0.987 (95% CI 0.127; 1.846), $p = .024$; group-based therapy 0.953 (95% CI 0.185; 1.722), $p = .015$; psychotherapy 0.607 (95% CI 0.147; 1.066), $p = .010$; collaborative care 0.207 (95% CI 0.050; 0.364), $p = .010$. Effect sizes for exercise ($p = .121$) online treatment ($p = .499$) and phone treatment ($p = .830$) were not significant. The forest plot is shown in Figure 3.

Eight studies in patients with diabetes and subthreshold depressive symptoms (Ebert et al., 2017; Hermanns et al., 2015; Huang et al., 2016; Penckofer et al., 2012; Pibernik-Okanovic et al., 2009, 2015; Tovote et al., 2014; Zheng et al., 2015) improved depression outcomes with an effect size of 0.360 (95% CI 0.204; 0.516), $p < .0001$ for all treatments. For psychotherapy: 1,131 (95% CI

0.083; 2.178), $p = .034$; and for online treatment 0.737 (95% CI 0.484; 0.990), $p < .0001$. Group therapy and psychoeducation had no significant effect. Glycemic control outcome effect sizes were significant for psychotherapy: 0.927 (95% CI 0.399; 1.455), $p = .001$ and group therapy: 0.237 (95% CI 0.019; 0.454), $p = .033$ (Appendix pp 13–14).

The meta-regression analysis showed a significant association (slope 0.137; $p < .0001$) between baseline HbA_{1c} and HbA_{1c} as outcome but no association with depression as outcome. High baseline HbA_{1c} was associated with a greater reduction in HbA_{1c}. There was a significant association (slope of 0.023; $p = .018$) between severity of depression at baseline and depression as treatment outcome; and between severity of depression at baseline and glycemic control as outcome (slope 0.028; $p = .005$). High baseline depression score was associated with a greater reduction in HbA_{1c} and depressive outcome (Appendix pp 15–18).

I^2 values for the pooled outcomes were of moderate heterogeneity (Higgins et al., 2019) (69%) for all outcomes combined in all included studies. Based on the residuals, there were no outliers. This indicates that there is a distribution of intervention effects, as was expected as different interventions were compared. A fixed model meta-analysis performed in all studies and outcomes as a meaningful test of the null hypothesis that there is no effect in every study (Higgins et al., 2019) showed p -value $< .0001$ (Appendix pp. 19) indicating that the interventions were effective. Irrespective of the scales used and outcomes measured, consistent beneficial effects were

TABLE 1 Data extraction table (N = 43)

Author (year)	n (completers) Mean age	Measure for depression classification	Intervention conditions and follow-up	Baseline depression, diabetes (Mean, SD)
Psychotherapeutic interventions (11 RCTs, n = 1,010)				
Lustman et al. (1998) USA	n = 41 Type 2 DM 100% 53.1- 56.4 ± 10.5-9.7	MDD according to DIS and BDI ≥ 14	CBT plus diabetes education versus diabetes education alone (CAU) FU: 11 wk, 6 months	Depression: BDI: CBT = 24.9 (10.2); Control = 21.1(6.8). Diabetes: HbA _{1c} : CBT = 10.2(3.6); Control = 10.4 (3.1)
Huang et al. (2002) China	n = 59 Type 2 DM 100%	SDS > 50	Antidiabetics + diabetic education + psychological treatment + relaxation and music treatment versus. Antidiabetics only (CAU) FU: 3 months	n/a
Li et al. (2003) China	n = 120 Type DM % not stated 50.5- 52.3 ± 10.4- 11.2	SDS ≥ 50	Antidiabetics + diabetic education + psychological treatment versus. antidiabetics only (CAU) FU: 4 wk	n/a
Lu et al. (2005) China	n = 60 Type 2 DM 100% 65.6-64.9 ±9.8-9.5	Mental maladjustment caused by CVA according to the CCMD-2-R and HAMD-17 ≥ 8	Diabetes and CVA education + electromyographic treatment + psychological treatment versus. CAU FU: 4 wk	Depression: HAMD: study group = 16.2 (5.7) control group = 16.8 (5.1). FBG: study group = 9.76 (3.83); control group = 9.89(3.94). Diabetes: 2HPG: study group = 13.65(4.72); control group = 13.31(4.57).
Simson et al. (2008) Germany	n = 30 Type 1 and Type 2 DM 80% 60.5 (±10.9)	HADS depression score ≥ 8	Individual supportive psychotherapy versus. CAU FU: discharge (3- 20 wk)	Depression: HADS-D: Psychotherapy = 11.7 (2.7); Control = 10.6 (2.9). Diabetes: HbA _{1c} : Psychotherapy = 7.8 (1.5); Control = 8.7 (1.8).
Piette et al. (2011) USA	n = 291 Type 2 DM = 100% 56.0 (±10.1)	BDI ≥ 14	Telephone delivered CBT plus walking program versus Enhanced CAU FU: 12 months.	Depression: BDI: EUC = 26.5 (9.9); CBT = 26.7 (7.7) Diabetes: HbA _{1c} : EUC = 7.7(1.7); CBT = 7.5 (1.7).
Safren et al. (2014) [89] USA	n = 78 Type 2 DM = 100% 55.44-58.31 (±8.72-7.41)	MDD as defined by DSM-IV assessed by clinician using MINI	CBT for adherence and depression versus enhanced CAU FU: 4, 8 and 12 months	Depression: MADRS: CBT-AD = 25.6(8.99); ETAU = 23.31(7.20). Diabetes: HbA _{1c} : CBT-AD = 8.81(1.78); ETAU = 8.74(1.41).
Tovote et al. (2014) Netherlands	n = 91 Type 2 DM = 61% Mean age = 53.1 (±11.8)	BDI-II ≥ 14	8-weekly sessions of Mindfulness based cognitive therapy versus CBT versus wait list control. FU: 3 months	Depression: BDI-II: MBCT = 23.6(7.7); CBT = 25.6(8.7); control = 24.3(8.0); HAM-D7: MBCT = 8.9(3.5); CBT = 9.4 (3.8); control = 7.5(2.8). Diabetes: HbA _{1c} : MBCT = 8.0(0.9); CBT = 8.3(1.4)
Schneider et al. (2016) USA	n = 29 Type 2 DM = 100% 53.4 (±7.1) 100% female	MDD as defined by SCID-IV	Behavioral action with exercise versus enhanced CAU. FU: 3 and 6 months	Depression: BDI-II: EUC = 21.6 (4.7); EX = 18.5 (8.2); HDSR: EUC = 17.4 (4.3); EX = 15.7(4.6) Diabetes: HbA _{1c} : EUC = 7.9(0.6); EX = 7.9 (0.8).

(Continues)

TABLE 1 (Continued)

Outcome assessment; depression, diabetes	Effect size depression, diabetes	Comments	MDD/sub threshold	Add-on exercise	Adherence focus
<i>Depression:</i> Response (reduction BDI \geq 50%) $p < .001$ in CBT group <i>Diabetes:</i> HbA _{1c} lower in CBT group $p < .03$	<i>Depression:</i> $\Delta - 1.112$ <i>Diabetes:</i> $\Delta - 0.704$	Improvement in depression as well as glycemic control in CBT versus. control	MDD	No	No
<i>Depression:</i> SDS total score difference in means 0.07; $p < .05$ <i>Diabetes:</i> HbA _{1c} difference in means 1.7; $p < .05$	<i>Depression:</i> $\Delta - 0.521$ <i>Diabetes:</i> $\Delta - 0.521$	Improvement in depression as well as glycemic control in CBT versus. control.	MDD	No	No
<i>Depression:</i> SDS total score difference in means 13.4, $p < .01$ <i>Diabetes:</i> FBG difference means 2.09, $p < .05$	<i>Depression</i> $\Delta - 0.478$: <i>Diabetes:</i> $\Delta - 0.362$	Anxiety (SAS \geq 50) taken into account as well. Improvement in depression as well as glycemic control in CBT versus. control	MDD	No	No
<i>Depression:</i> HAMD-17 total score difference in means 7.3; $p < .01$ <i>Diabetes:</i> difference in means FPG 1.54; $p < .05$	<i>Depression</i> $\Delta - 0.688$: <i>Diabetes:</i> $\Delta - 0.517$	Hemiplegia after CVA as DM complication. Improvement in depression as well as glycemic control in CBT versus. control	MDD	No	No
<i>Depression:</i> HADS depression scale total score mean difference 1.9; $p = .018$ <i>Diabetes:</i> PAID mean difference 7.6; $p = .008$	<i>Depression:</i> $\Delta - 0.918$ <i>Diabetes:</i> $\Delta - 1.043$	Diabetic foot as DM complication; Inpatients. Improvement in depression as well as glycemic control in supportive psychotherapy versus. control	MDD	No	No
<i>Depression:</i> BDI total score mean between group difference -4.5, $p < .0001$ <i>Diabetes:</i> HbA _{1c} mean between group difference 0.07, $p = .70$.	<i>Depression:</i> $\Delta 0.418$ <i>Diabetes:</i> $\Delta 0.000$	Improvement in depressive symptoms but not glycemic control for telephone CBT + walking versus. control	MDD	Yes	No
<i>Depression:</i> MADRS score mean difference 6.22 ($p = .002$). CGI ratings mean difference 0.74, ($p = .01$) <i>Diabetes:</i> HbA _{1c} mean difference 0.72, $p = .001$.	<i>Depression:</i> $\Delta 0.762$ <i>Diabetes:</i> $\Delta 2.311$	Main focus on adherence. Significant improvements in depression as well as glycemic control in CBT-AD versus. control.	MDD	No	Yes
<i>Depression:</i> BDI-II scores and HAM-D7 showed significant improvements in both interventions ($p < .001$). There was a clinically relevant improvement of 26% (MBCT) and 29% (CBT) versus. 4% (control). <i>Diabetes:</i> HbA _{1c} levels did not change significantly after MBCT ($p = .92$) or CBT ($p = .72$)	MBCT: <i>Depression:</i> $\Delta 0.568$ <i>Diabetes:</i> n/a CBT: <i>Depression:</i> $\Delta 0.541$ <i>Diabetes:</i> n/a	Significant improvement in depressive symptoms for both MBCT and CBT versus wait list control. HbA _{1c} levels did not improve in either intervention group.	Sub	No	No
<i>Depression:</i> BDI-II total score mean difference -7.3, $p < .0001$, HRSD mean difference score = -6.6, ($p < .0001$). <i>Diabetes:</i> Time x condition was not significant for HbA _{1c} ($p = .78$).	<i>Depression:</i> $\Delta - 0.018$ <i>Diabetes:</i> $\Delta - 0.114$	Significant improvement in depressive symptoms in both EX group and EUC group. No improvement in glycemic control.	MDD	Yes	No

(Continues)

TABLE 1 (Continued)

Author (year)	n (completers) Mean age	Measure for depression classification	Intervention conditions and follow-up	Baseline depression, diabetes (Mean, SD)
Huang et al. (2016) China	n = 61 Type 2 DM = 100% 56.43 (±10.42)	CES-D=>16 (indicating "significant" or mild depressive symptomatology)	Combined behavioral intervention 12 sessions over 3 months versus CAU FU: 3 months	Depression: CES-D: CBT + MET = 21.81 (5.68); EUC = 21.97 (3.37). Diabetes: HbA _{1c} : CBT + MET = 7.68(1.44); EUC = 7.84 (1.95).
De Groot et al. (2019) USA	n = 140 Type 2 DM = 100% 56.0 (±10.7)	Structured clinical interview for DSM_IV-TR, BDI-II	CBT alone versus community-based exercise versus CBT + exercise versus CAU FU: 12 weeks	Depression: BDI-II: (values not reported) Diabetes: HbA _{1c} : CBT = 8.0(1.6); EX = 8.1(1.7); CBT + EX=7.5(1.6); UC = 8.0(1.9).
Comparison of different pharmacological agents (6 RCTs, comparative effectiveness trials (CETs) n = 291)				
Gülseren et al. (2005) Turkey	n = 23 Type 2 DM 100% 58.2-57.1 ± 12.3-10.4	MDD according to SCID, HADS depression subscale score ≥ 10, HDRS ≥ 16	Fluoxetine versus. paroxetine FU: 12 wk	Depression: HDRS: Fluoxetine = 17.5 (2.4); Paroxetine = 18.8(3.0). Diabetes: HbA _{1c} : Fluoxetine = 6.9(1.7); Paroxetine = 6.9(1.2)
Barragan-Rodrigues et al. (2008) Mexico	n = 23 Type 2 DM = 100%, 66.4-69 (±5.9-6.1)	Ysavage and Brink score > 11.	Magnesium supplementation versus imipramine FU: 12 weeks	Depression: Ysavage & Brink: MgCl ₂ = 17.9(3.9); Imipramine = 16.1(4.5). Diabetes: HbA _{1c} : MgCl ₂ = 8.9(1.6) Imipramine = 9.0(1.7); FBG: MgCl ₂ = 194.3 (59.0); Imipramine = 183.4(68.0).
Khazaie et al. (2011) Iran	n = 47 Type 2 DM = 100% 47.7-51.57 (±8.63-8.38)	BDI ≥ 14	Fluoxetine versus Citalopram FU:12 weeks	Depression: BDI: Fluoxetine = 29.29 (3.50); Citalopram = 25.26(3.51). Diabetes: HbA _{1c} : Fluoxetine = 7.68 (1.69); Citalopram = 8.25 (1.34); FBG: Fluoxetine = 159.21 (39.66); Citalopram = 163.37 (49.24).
Karaiskos et al. (2013) Greece	n = 40 Type 2 DM = 100% 52.4-54.3 (±11.4-12.5)	Classification of mood disorder based on DSM-IV-TR criteria	Agomelatine versus Sertraline FU:4 months	Depression: HDRS: Agomelatine = 11.6(2.5); Sertraline = 11.5(2.5). Diabetes: HbA _{1c} : Agomelatine = 7.7(0.5); Sertraline = 7.6(0.5); FBG: Agomelatine = 137(21); Sertraline = 135(16).
Kang et al. (2015) China	n = 116 Type 2 DM = 100% 50.82-52.50 (±11.36-10.27)	HDRS ≥ 17 (indicating moderate-to-severe depression) + psychiatrist's assessment according to DSM criteria.	Agomelatine versus paroxetine FU: 6,12 weeks	Depression: HDRS: Paroxetine = 23.94(3.07) Agomelatine = 24.20(3.38) Diabetes: HbA _{1c} : Paroxetine = 7.71, Agomelatine = 7.84 (0.45).

(Continues)

TABLE 1 (Continued)

Outcome assessment; depression, diabetes	Effect size depression, diabetes	Comments	MDD/sub threshold	Add-on exercise	Adherence focus
<p><i>Depression:</i> CES-D scores reduced significantly ($-6.28, p < .01$) in intervention group.</p> <p><i>Diabetes:</i> HbA_{1c} levels reduced significantly in the intervention group (-4.86) and were significantly lower in the intervention group than control ($p < .01$).</p>	<p><i>Depression:</i> Δ 2.298</p> <p><i>Diabetes:</i> Δ 0.915</p>	Improvement in depressive symptoms and glycemic control for MET + CBT versus. control	Sub	No	No
<p><i>Depression:</i> Full remission from MDD was 66% in CBT, 72% in Ex, 71% in CBT + Ex compared with 32% in CAU. BDI-II scores lower in all three groups compared to CAU ($p < 0.05$).</p> <p><i>Diabetes:</i> HbA_{1c} levels lower for CBT + ex ($p = .0016$), but not CBT ($p = .379$) or Ex alone ($p = .132$)</p>	<p>CBT: <i>Depression:</i> Δ 0.678</p> <p><i>Diabetes:</i> Δ 0.274;</p> <p>Exercise: <i>Depression:</i> Δ 0.640;</p> <p><i>Diabetes:</i> Δ 0.467;</p> <p>CBT + Ex: <i>Depression:</i> Δ 0.671;</p> <p><i>Diabetes:</i> Δ 0.912.</p>	Significantly larger improvement in depressive symptoms in CBT alone, exercise alone and CBT + exercise compared with CAU. Glycemic control only showed significant improvements in CBT + exercise group versus CAU.	MDD	Yes	No
<p><i>Depression:</i> Both groups improved significantly in HDRS scores (mean difference 0.62; $p = .003$)</p> <p><i>Diabetes:</i> No difference in HbA_{1c} (mean diff 0.11; n.s.)</p>		No significant difference between both conditions. This study is not pooled in the meta-analysis.	MDD	No	No
<p><i>Depression:</i> Ysavage and brink scores reduced for MgCl₂ ($-6.5; p < .005$) and imipramine ($-6.2; p < .005$).</p> <p><i>Diabetes:</i> No change for HbA_{1c} or FBG levels.</p>	n/a	Depressive symptoms but not glycemic control improved in both MgCl ₂ and imipramine groups. No control group so study not included in meta-analysis.	MDD	No	No
<p><i>Depression:</i> BDI scores improved for fluoxetine ($-15.12, p < .001$) and citalopram ($11.84, p < .001$).</p> <p><i>Diabetes:</i> HbA_{1c} levels improved for fluoxetine ($-1.94; p < .001$) and citalopram ($-1.59; p < .001$). FBG also improved in Fluoxetine ($48.93; p < .001$) and citalopram ($39.95; p < .001$).</p>	n/a	Fluoxetine and Citalopram both improved depression symptoms and glycemic control. No control group so study not included in meta-analysis.	Sub	No	No
<p><i>Depression:</i> HDRS scores reduced more for Agomelatine (-5.8) than sertraline (-4.2) ($p = .050$).</p> <p><i>Diabetes:</i> No effect on FPG levels ($p = .694$). HbA_{1c} levels reduced more for Agomelatine (-0.5) than sertraline (-0.0) ($p = .044$).</p>	n/a	Depressive symptoms and HbA _{1c} levels but not FBG levels were significantly lower in Agomelatine group compared to the sertraline group at follow-up. No control group so study not included in meta-analysis.	MDD	No	No
<p><i>Depression:</i> HDRS scores improved for Agomelatine (-14.85) and paroxetine (-13.44) ($p < 0.001$).</p> <p><i>Diabetes:</i> HbA_{1c} levels significantly lower for Agomelatine ($\Delta -0.62, p < .001$) but not paroxetine ($p = .038$).</p>	n/a	Both drugs significantly improved depressive symptoms. Agomelatine better than paroxetine for glycemic control. No control group so study not included in meta-analysis.	MDD	No	No

(Continues)

TABLE 1 (Continued)

Author (year)	n (completers) Mean age	Measure for depression classification	Intervention conditions and follow-up	Baseline depression, diabetes (Mean, SD)
Kumar et al. (2015) India	n = 42 Type 2 DM = 100% 48.65–49.75 (±10.19–14.27)	HDRS ≥ 14	Agomelatine versus Escitalopram FU: 1 month, 2 months	<i>Depression:</i> HDRS: Escitalopram = 18.5 (2.95) Agomelatine = 17.15(2.54) MADRS: Escitalopram = 22.95(2.74) Agomelatine = 17.15(2.54) <i>Diabetes:</i> HbA _{1c} : Escitalopram = 7.2 (0.36) Agomelatine = 7.35 (0.41) FBG: Escitalopram = 131.9 (3.45) Agomelatine = 132.3 (4.41)
Pharmacological versus psychotherapeutic (2 RCTs, CETs, n = 149)				
Gois et al. (2014) Portugal	n = 34 Type 2 DM = 100% 55.14 (±5.92)	HADS > 7; MADRS > 17 & MDD according to MINI	Sertraline and clinical management versus Interpersonal psychotherapy FU: 6 weeks, 14 weeks, 24 weeks	<i>Depression:</i> MADRS: Sertraline = 24.64(6.4); IPT = 26.45 (4.37). <i>Diabetes:</i> HbA _{1c} : Sertraline = 9.28(2.67); IPT = 8.69(2.20).
Petrak, Baumeister, et al. (2015) Germany	n = 251 (n = 115) Type 2 DM = 48.6% 48.5 1(±1.7)	CES-D > 22	Diabetes specific CBT versus sertraline FU: 12 weeks (phase 1), 15 months (excluding nonremitting patients at phase 1)	<i>Depression:</i> HAMD-17: CBT = 18.04(4.62); Sertraline = 18.87(5.14); <i>Diabetes:</i> HbA _{1c} : CBT = 9.37(1.63); Sertraline = 9.15(1.37).
Pharmacological treatment versus placebo (7 RCTs, n = 339)				
Lustman et al. (1997) USA	n = 28 Type 1 and Type 2 DM 50% 49.0–49.2 ± 12.1–13.7	MDD according to DIS	Glucometer training + nortriptyline versus placebo FU: 9 wk	<i>Depression:</i> BDI: Nortriptyline = 19.0 (7.4); Placebo = 17.8 (7.1) <i>Diabetes:</i> HbA _{1c} : Nortriptyline = 11.8 (2.9) Placebo = 11.6 (3.1) (*Depressed group n = 28)
Lustman, Anderson, et al. (2000) USA	n = 54 Type 1 and Type 2 DM 55.6% 45.0–47.7±13.0–11.5	MDD (DIS), and BDI or HAMD ≥ 14	Fluoxetine versus placebo FU: 8 wk	<i>Depression:</i> BDI: Fluoxetine = 23.6 (8.2); Placebo = 22.4 (9.1); HAMD: Fluoxetine = 20.1 (5.6); Placebo = 19.5 (6.9). <i>Diabetes:</i> HbA _{1c} : fluoxetine = 8.4 (1.7); Placebo = 8.6 (1.6).
Paille-Hyvärinen et al. (2003) Finland	n = 13 Type 2 DM 100% 61.1–62.3 ±8.6–11.5	MADRS score between 2.5 and 12 (mild-to-moderate depression)	Paroxetine versus placebo FU: 4 wk	<i>Depression:</i> MADRS: Placebo = 6.4 (4.0); Paroxetine = 7.4(2.9); BDI: Placebo = 13.0 (9.2); Paroxetine = 13.7 (7.4). <i>Diabetes:</i> HbA _{1c} : Placebo = 6.9 (0.4); Paroxetine = 7.5(0.8)
Xue (2004) China	n = 48 Type 1 and Type 2 DM 85.4% 21–65 age range		Paroxetine versus placebo FU: 8 wk	<i>Depression:</i> HAMD: Paroxetine = 20.1(12.7; control = 19.5 (12.1); BDI: Paroxetine = 23.6(14.2); control = 22.4(15.9). <i>Diabetes:</i> GHb: paroxetine = 8.8(1.8); control = 8.7(1.6).

(Continues)

TABLE 1 (Continued)

Outcome assessment; depression, diabetes	Effect size depression, diabetes	Comments	MDD/sub threshold	Add-on exercise	Adherence focus
<i>Depression:</i> Larger improvement in Escitalopram group for HDRS scores (-9.65 v -1.55, $p < .001$); and MADRS scores (-9.35 v -2.00, $p < .001$) <i>Diabetes:</i> Larger HbA _{1c} improvement in Escitalopram (-0.09) than Agomelatine (-0.03) ($p = .047$). Larger reduction in FBG in Escitalopram (-6.95) than Agomelatine (-4.45) ($p = .043$).	n/a	Escitalopram appears to be better than Agomelatine for improving both depression and glycemic control. No control group so study not included in meta-analysis.	Sub	No	No
<i>Depression:</i> MADRS scores improved in both IPT (-14.00) and sertraline (-14.00) ($p < 0.0001$). <i>Diabetes:</i> No significant improvement in HbA _{1c} levels.	n/a	No differences in improvements in depressive symptoms between IPT and sertraline. No significant effect on glycemic control was shown for either intervention.	MDD	No	No
<i>Depression:</i> HAMD-17 scores improved in CBT (-10.21) and sertraline (-13.41). Sertraline improvement was significantly better ($p > .05$). <i>Diabetes:</i> No significant change in HbA _{1c} levels.	n/a	Sertraline and CBT both improve depression after 12 weeks. Significant advantage of sertraline over diabetes specific CBT for improving depressive symptoms over one year, but not glycemic control.	MDD	No	No
<i>Depression:</i> BDI total score, mean difference 5.6; $p = .03$ <i>Diabetes:</i> HbA _{1c} , no significant difference, no outcome reported.	<i>Depression:</i> $\Delta - 0.868$ <i>Diabetes:</i> $\Delta 0$	Poorly controlled (HbA _{1c} $\geq 9\%$) as inclusion criterion. Improvement in depression but not in glycemic control in nortriptyline versus. control. Nortriptyline may have negative impact on glycemic control.	MDD	No	No
<i>Depression:</i> HAMD total score mean difference 26.7; $p < .04$ <i>Diabetes:</i> HbA _{1c} mean difference 0.33; $p = .13$ (n.s.)	<i>Depression:</i> $\Delta - 0.573$ <i>Diabetes:</i> $\Delta 0.419$	Improvement in depression but not in glycemic control in fluoxetine versus. placebo.	MDD	No	No
After initial improvement in Paroxetine group at 3 months, no significant improvement for both outcomes at end of follow-up. <i>Depression:</i> MADRS total score mean difference 2.50; $p = .25$ (n.s.) <i>Diabetes:</i> GHbA _{1c} mean difference 0.37; $p = .08$ (n.s.)	<i>Depression:</i> $\Delta - 0.676$ <i>Diabetes:</i> $\Delta 1.073$	Poorly controlled (HbA _{1c} $\geq 6.5\%$ or FBG ≥ 7.0) as inclusion criterion. Probably a combination of ceiling effect and underpowered study.	MDD	No	No
<i>Depression:</i> HAMD-17 total score mean difference 5.7; $p < .01$ <i>Diabetes:</i> HbA _{1c} mean difference 0.4; $p = .245$ (n.s.)	<i>Depression:</i> $\Delta - 0.776$ <i>Diabetes:</i> $\Delta 0.340$	Improvement in depression but not in glycemic control in paroxetine versus. placebo.	MDD	No	No

(Continues)

TABLE 1 (Continued)

Author (year)	n (completers) Mean age	Measure for depression classification	Intervention conditions and follow-up	Baseline depression, diabetes (Mean, SD)
Paile-Hyvärinen et al. (2007) Finland	n = 49 Type 2 DM 100% 59.5–59.2 ± 6.0–5.4	Diagnostic interview. Mild depression (< 6 depressive symptoms according to the DSM-IV).	Paroxetine versus placebo FU: 3 months, 6 months	Depression: HADS-D: Placebo = 8.4 (3.4); Paroxetine = 7.3 (3.4). Diabetes: Placebo = 8.7(1.3); Paroxetine = 8.5 (0.9).
Echeverry et al. (2009) USA	n = 89 n = 87 Type 2 DM; n = 2 Type 1 52–53 ±8–10	MDD according to CDIS	Sertraline versus placebo	Depression: HAM-D: Sertraline = 19.0(5.0); Control = 20.0 (6.0); Diabetes: HbA _{1c} : Sertraline = 10.0(1.8) Control = 9.7(1.6).
Guo et al. (2014) China	n = 58 Type 2 DM = 100% 53.3–54.7(±7.3–7.3)	Classification of depression based on DSM-IV criteria	Metformin versus placebo FU: 24 weeks	Depression: MADRS: Metformin = 23.7 (3.5), placebo = 24.3 (3.8); HRSD17: Metformin = 20.1 (3.0); placebo = 20.4 (2.4). Diabetes: HbA _{1c} : Metformin = 7.82(0.82); placebo = 8.01(0.59).
Psychoeducation (2 RCTs, n = 259)				
Pibernik-Okanovic et al. (2009) Croatia	n = 50 Type 2 DM = 100% Median age = 55 (51–62)- 58 (53–64)	Mild-to-moderate depression. PHQ9 scores 10–14	Psychoeducation comprising 4 x interactive group meetings versus CAU (Depression screening followed by standard diabetes treatment) FU: 6, 12 months	Depression: CES-D: PsyEd = 26(22–30); CAU = 24 (18–35). Diabetes: HbA _{1c} : PsyEd = 7.5(6.4–8.3); CAU = 7.7(6.6–8.9). *medians (CI)
Pibernik-Okanovic et al. (2015) Croatia	n = 209 Type 2 DM = 100% 57.7–58.5 (±6.2–5.6)	PHQ2 - > one depressive symptom over last month	Psychoeducation versus psychoeducation and physical exercise) versus enhanced CAU FU: 12 months	Depression: CES-D: PsyEd = 19.7(9.1); PsyEd + Ex =19.8(8.2); CAU = 19.0(8.6). Diabetes: HbA _{1c} : PsyEd = 7.4(1.3); PsyEd + Ex =7.2(1.0); CAU = 7.1(1.0).
Collaborative Care (6 RCTs, n = 1,133)				
Katon, Von Korf, et al. (2004) USA	n = 329 Type 1 and Type 2 DM 95.7% 58.1–58.6 ± 12.0–11.8	PHQ-9 ≥ 10 and SCL-90 depression mean item score > 1.1	Collaborative care versus CAU FU: 6 months, 12 months	
Williams et al. (2004)USA	n = 417 Type 1 and Type 2 DM % not stated, mostly Type 2 71.2 ± 7.5	MDD or dysthymia according to SCID	Education about late-life depression + collaborative care versus CAU FU: 3 months, 6 months, 12 months	
Ell et al. (2010) USA	n = 387 Type 1 and Type 2 DM % group not stated 72% >50 years	PHQ-9 ≥ 10	Collaborative care versus CAU FU: 6 months, 12 months, 18 months	

(Continues)

TABLE 1 (Continued)

Outcome assessment; depression, diabetes	Effect size depression, diabetes	Comments	MDD/sub threshold	Add-on exercise	Adherence focus
<i>Depression:</i> HADS depression scale total score mean difference 0.7; $p=.448$ (n.s.). <i>Diabetes:</i> GHbA _{1c} mean difference 0.13; $p=.693$ (n.s.).	<i>Depression:</i> $\Delta - 0.260$ <i>Diabetes:</i> $\Delta 0.135$	No significant improvement in depressive outcomes and glycemic control.	MDD	No	No
<i>Depression:</i> HADS depression scale total score mean difference 1.0; (n.s.). <i>Diabetes:</i> GHbA _{1c} mean difference 1.1; $p<.011.0$; (n.s.).	<i>Depression:</i> $\Delta - 0.283$ <i>Diabetes:</i> $\Delta - 0.480$	Significant improvement in depression in both sertraline and placebo; no difference between conditions. Significant improvement in glycemic control in sertraline compared to placebo.	MDD	No	No
<i>Depression:</i> MADRS ($p<.001$) and HRSD-17 ($p<.001$) scores both improved for Metformin <i>Diabetes:</i> HbA _{1c} levels improved compared to placebo group (-1.52 versus 0.19 $p<.001$).	<i>Depression:</i> $\Delta 0.900$ <i>Diabetes:</i> $\Delta 3.676$	Significant improvement in depressive symptoms and glycemic control in metformin group compared to placebo group.	MDD	No	No
<i>Depression:</i> Median CES-D scores reduced in both groups. Between group difference n.s ($p=.074$) <i>Diabetes:</i> HbA _{1c} levels reduced, between group difference n.s ($p=.089$).	<i>Depression:</i> $\Delta 0.135$ <i>Diabetes:</i> $\Delta -0.049$	Psychoeducation shows no significant benefit for either depressive symptoms or glycemic control over care as usual.	Sub	No	No
<i>Depression:</i> CES-D improved for all groups ($p=.003$) but not significantly between groups ($p = .656$) <i>Diabetes:</i> No significant effects on HbA _{1c} levels	PsyEd: <i>Depression:</i> $\Delta 0.082$ <i>Diabetes:</i> $\Delta -0.210$. PsyEd + Ex: <i>Depression:</i> $\Delta -0.074$ <i>Diabetes:</i> $\Delta -0.199$	Psychoeducation and psychoeducation + exercise showed no significant benefits over treatment as usual for depressive symptoms or glycemic control.	Sub	Yes	No
<i>Depression:</i> SCL-20 total score mean difference response (reduction SCL-90 $\geq 40\%$ or $\geq 50\%$) $p=.004$ <i>Diabetes:</i> HbA _{1c} mean difference 0, n.s.	<i>Depression:</i> $\Delta - 0.320$ <i>Diabetes:</i> $\Delta 0.085$	Improvement in depression but not in glycemic control in collaborative care versus. usual care.	MDD	No	No
<i>Depression:</i> SCL-20 total score mean difference - 0.3; CI - 0.57 to 0.29 <i>Diabetes:</i> HbA _{1c} mean difference 0, n.s.	<i>Depression:</i> $\Delta - 0.676$ <i>Diabetes:</i> $\Delta 0.000$	Improvement in depression but not in glycemic control in collaborative care versus. usual care.	MDD	No	No
<i>Depression:</i> SCL-20 total score mean difference 50% improved in 62 versus. 44%; $p<.001$ <i>Diabetes:</i> HbA _{1c} mean difference 0, n.s.	<i>Depression:</i> $\Delta - 0.337$ <i>Diabetes:</i> $\Delta - 0.263$	Significant improvement in depression but not in glycemic control in collaborative care versus. usual care in Hispanics with baseline HbA _{1c} > 8	MDD	No	No

(Continues)

TABLE 1 (Continued)

Author (year)	n (completers) Mean age	Measure for depression classification	Intervention conditions and follow-up	Baseline depression, diabetes (Mean, SD)
Ell et al. (2011) USA	n = 387 Type 2 DM = 98% 54 (±8.7)	PHQ9 scores > 10	Socioculturally adapted collaborative care (MDDP; n = 193) versus enhanced CAU (EUC: n = 194). FU: 6, 12, 18, 24 months	Depression: SCL-20; PHQ9 (values not reported). Diabetes: HbA _{1c} (values not reported).
Bogner et al. (2012) USA	n = 180 Type 2 DM = 100% 57.1-57.8 (9.6-9.4)	PHQ9	Integrated care versus CAU FU: 6 and 12 weeks	Depression: PHQ9: IC = 10.6(7.9); CAU = 9.9(7.2). Diabetes: HbA _{1c} : IC = 7.2(1.8); CAU = 7.0(1.9).
Johnson et al. (2014) [28] USA	n = 157 Type 2 DM = 100% 57.0-59.2 (±10.5-8.5)	PHQ scores > 10	TEAMCare (n = 95) collaborative care intervention versus screening and follow-up CAU (control; n = 62). FU: 6 months, 12 months	Depression: PHQ9: TEAMCare = 14.5(3.8); Control = 14.6(3.5). Diabetes: HbA _{1c} : TEAMcare = 7.5(1.8); Control = 7.8(1.7).
Online-based interventions (3 RCTs, n = 605)				
Van Bastelaar et al., (2011) Netherlands	n = 255 Type 2 DM = 55% 50 (±12)	CES-D > 16	Web-based CBT (iCBT; n = 125) versus waiting list control (WL; n = 130) group. FU: 1 month	Depression: CES-D: iCBT = 29(7); WL = 28(7); Diabetes: HbA _{1c} : iCBT = 7.4(1.6); WL = 7.3(1.4).
Ebert et al. (2017) [79] Germany	n = 260 Type 2 DM = 55% 50.8 (±11.8)	CES-D > 23	GET ON. Mood Enhancer Diabetes - Internet guided self-help intervention (n = 129) for depression versus CAU + online education on depression FU: 8 weeks, 6 months	Depression: HADS-D: GET ON = 12.0(3.2); CAU = 11.7(3.7). Diabetes: HbA _{1c} : GET ON = 7.6(1.6); CAU = 7.4(1.3).
Newby et al. (2017) Australia	n = 90, Type 2 DM = 42%, 46.7 (±12.6)	PHQ9 scores 5=>23	Web-based CBT versus CAU FU: 3 months (for iCBT group only)	Depression: PHQ9: iCBT = 15.95(5.25); TAU = 14.29(5.25). Diabetes: HbA _{1c} : iCBT = 7.87(1.79); TAU = 7.72(1.82).
Group-based interventions (4 RCTs, n = 508)				
Penckofer et al. (2012) USA	n = 74 Type 2 DM = 100% Female = 100% 54.0-54.8 (±8.4-8.8)	>16 CES-D (indicating "significant" or mild depressive symptomatology - average of 2 screenings)	SWEEP psychoeducational intervention versus CAU FU: 3 and 6 months.	Depression: CES-D: SWEEP = 27.7(9.3); UC = 28.9(9.5). Diabetes: HbA _{1c} : SWEEP = 7.8(1.8); UC = 7.9(2.0); FBG: SWEEP = 165.3 (71.1); UC = 168.8 (74.9).

(Continues)

TABLE 1 (Continued)

Outcome assessment; depression, diabetes	Effect size depression, diabetes	Comments	MDD/sub threshold	Add-on exercise	Adherence focus
<p>Depression: SCL-20 and PHQ 9 scores improved significantly more in intervention group ($p < 0.001$).</p> <p>Diabetes: No differences in HbA_{1c} levels ($p > 0.05$).</p>	n/a	<p>Significantly larger improvements in depressive symptoms were observed in the MDDP group versus care as usual, however these group differences narrowed over time. No effects on glycemic control.</p> <p>Study not entered in meta-analysis due to lack of data.</p>	MDD	No	No
<p>Depression: PHQ-9 scores improved significantly more in IC group (-2.42; $p = .007$). IC group were more likely to achieve remission (58.7% versus 30.7%; $p < .001$)</p> <p>Diabetes: HbA_{1c} levels significantly improved in IC group (-0.70 $p < .001$).</p>	<p>Depression: Δ 0.405</p> <p>Diabetes: Δ 0.497</p>	Improvement in glucose control and depressive symptoms in integrated care intervention versus usual care.	MDD	No	Yes
<p>Depression: PHQ9 scores improved significantly more in TEAMcare group (-7.3. ($p = .015$).</p> <p>Diabetes: No differences in HbA_{1c} levels</p>	<p>Depression: Δ 0.388</p> <p>Diabetes: Δ 0.244</p>	Significant improvement in depressive symptoms but not glycemic control in collaborative care group versus active control.	MDD	No	No
<p>Depression: Treatment x time interaction effect on CES-D scores ($p < .001$) was significant.</p> <p>Diabetes: No significant treatment effect found for HbA_{1c} levels ($p > .05$).</p>	n/a	Significant improvement in depressive symptoms but not glycemic control in web-based- CBT group versus active control. Study not entered in meta-analysis due to lack of reported data.	MDD	No	No
<p>Depression: CES-D total mean score difference = -7.7 ($p < .001$). HADs total mean score difference = -3.2 ($p < .001$).</p> <p>Diabetes: HbA_{1c} mean difference 0, n.s.</p>	<p>Depression: Δ 0.735</p> <p>Diabetes: Δ 0.133</p>	Significantly greater improvement in depressive symptoms in internet guided self-help versus active control. No effect on glycemic control.	Sub	No	No
<p>Depression: PHQ9 scores improved overall and the group x time interaction was significant ($p < .001$). 51% in iCBT versus 18% in TAU improved reliably.</p> <p>Diabetes: No significant interaction effect for HbA_{1c} levels ($p = .750$).</p>	<p>Depression: Δ 0.782</p> <p>Diabetes: Δ 0.142</p>	Significantly greater improvement in depressive symptoms but not glycemic control in Web-based CBT group versus care as usual. No follow-up data for care as usual group limits conclusions.	MDD	No	No
<p>Depression: CES-D scores mean difference = - 6.8 ($p < .01$).</p> <p>At 6 months 35% of intervention versus 80% of control remained depressed.</p> <p>Diabetes: No significant improvements for FBG or HbA_{1c} levels.</p>	<p>Depression: Δ 0.964</p> <p>Diabetes: Δ 0.272</p>	Significant improvement in depressive symptoms but not glycemic control in SWEEP psychoeducation group compared to control group.	Sub	No	No

(Continues)

TABLE 1 (Continued)

Author (year)	n (completers) Mean age	Measure for depression classification	Intervention conditions and follow-up	Baseline depression, diabetes (Mean, SD)
Hermanns et al. (2015) Germany	n = 214 Type 2 DM = 34.1% 43.3 (±13.3)	CES-D > 16	Self-management- orientated group program (DIAMOS) versus control group CAU + diabetes education. FU: 6 months, 12months	Depression: CES-D: DIAMOS = 24.4(7.5); CG = 22.1(8.6); HADS: DIAMOS = 10.9(4.3); CG = 9.6(3.8). Diabetes: HbA _{1c} : DIAMOS = 8.8(1.7); CG = 8.7(1.7).
Long et al. (2015) China	n = 100 Type 2 DM = 100% 66.8 (±9.03)	SDS > 50	8 sessions of group counseling versus CAU FU: 3, 6 and 12 months.	Depression: SDS: GC = 0.57(0.067); CAU = 0.58(0.055). Diabetes: HbA _{1c} : GC = 8.08(1.03); CAU = 8.10(1.10); FBG: GC = 9.26(1.70); CAU = 9.11(1.65).
Zheng et al. (2015) [92] China	n = 120 Type 2 DM = 100% 61-62 (±7-6)	Depression according to SDS	24 weeks Twenty-four move Shadow Boxing and psychosomatic relaxation versus control group with CAU community diabetes health instructions. FU:24 weeks.	Depression: SDS: Boxing = 53.2(8.5); Control = 54.3(9.2). Diabetes: HbA _{1c} : Boxing = 7.54(1.53); Control = 7.39(1.62).
Phone-based (1 RCT, n = 225)				
Naik et al., (2019) USA	n = 225 =61.9 (±8.3)	PHQ 9 scores > 10	Telehealth collaborative goal setting and behavioral activation versus enhanced CAU FU: 6 and 12 months.	Depression: PHQ9: HOPE = 15.8(4.2); EUC = 16.2(4.0); Diabetes: HbA _{1c} : HOPE = 9.2(1.4); EUC = 9.3(1.5).
Light Therapy (1 RCT, n = 83)				
Brouwer et al. (2019) Netherlands	n = 83 Type 2 DM = 100% 60.1-62.9 (±9.8-10.7)	IDS scores > 14 MDD according to DSM-IV criteria	Light therapy (active broad spectrum, white yellow light, 10,000 lux) versus placebo (monochromatic green light [545nm]) FU: 4, and 8 weeks	Depression: IDS: (values not reported). Diabetes: HbA _{1c} : Light = 7.2(1.1); Placebo = 7.2(1.3).

Note: The first column indicates the first author, year of publication and country study was conducted. The second column shows the sample size, % type 1 diabetes and type 2 diabetes and the Mean[SD] age of participants. The third column indicates how depressive disorder/presence of clinically significant symptoms or subthreshold disorder was diagnosed or defined. The fourth column describes the intervention, including the follow-up (FU) time periods. Column 5 shows the Baseline data for both diabetes (e.g., HbA_{1c}) and depression (e.g., depression questionnaire) outcomes. Column 6 shows the outcome data for both the diabetes and depression outcomes. Column 7 shows the effect size of the intervention on both the diabetes and depression outcomes. Column 8 describes the conclusions drawn from the study. Column 9 indicates whether the study focused on participants with depressive disorder or clinically significant symptoms (as noted by MDD) or subthreshold disorder (sub). Columns 10 and 11 show whether the intervention included an intervention component or focus on adherence, respectively. The number of trials and participants for each intervention is shown in the row indicating intervention type.

Abbreviations: BDI, beck depression inventory; CAU, care as usual; CBT, cognitive behavior therapy; CGI, clinical global impression; CVA, cerebrovascular accident; FBG, fasting blood glucose; HDRS, hamilton depression rating scale; MBCT, mindfulness-based cognitive therapy; MDD, major depressive disorder; SDS, self-rating depression scale.

findings from our meta-analysis enable us to tentatively propose a flowchart to guide treatment choice, based upon the clinical profile of the patient and building on existing guidelines for treatment of people with diabetes. This flowchart is shown in Figure 4.

Among the 32 randomized controlled trials included in the meta-analysis, most studies did not meet all criteria to reduce risk of bias, mostly due to unclear reporting about the method of randomization and blinding, instead the focus being on description of the intervention; unclear reporting about attrition rates and intention-to-treat

(ITT) analysis. Furthermore, one study had low rates of compliance with the intervention under study and unclear reporting about the numbers of compliant participants (Brouwer et al., 2019); ten studies used small underpowered samples, despite otherwise being of seemingly adequate quality. Details of the risk-of-bias assessment for included trials are provided in the appendix (pp 4-7). A sensitivity analysis in the 16 studies (Baumeister & Bengel, 2012; Begg, 1994; Beydoun & Wang, 2010; Biostat, 2005; Bot et al., 2010; DerSimonian & Laird, 1986; Guyatt et al., 2008; Higgins, 2003;

TABLE 1 (Continued)

Outcome assessment; depression, diabetes	Effect size depression, diabetes	Comments	MDD/sub threshold	Add-on exercise	Adherence focus
Depression: CES-D mean difference = -3.9 [95% CI 0.6-7.3] ($p = .021$) PHQ9 scores mean difference = -1.7 [95% CI 0.2-3.2] ($p = .023$). Diabetes: HbA _{1c} levels mean difference = -0.3, ($p = .230$)	Depression: $\Delta 0.039$ Diabetes: $\Delta 0.269$	Significant improvement in depressive symptoms but not glycemic control in DIAMOS group-based therapy versus control group.	MDD	No	No
Depression: SDS scores showed significant improvement ($p < .001$) Diabetes: FBG and HbA _{1c} levels showed significant difference between groups ($p < .05$)	Depression: $\Delta 1.637$ Diabetes: $\Delta 0.927$	Improvement in depression scores, fasting blood glucose and glycemic control in group counseling versus usual care	MDD	No	No
Depression: SDS scores mean difference = -4.0 ($p < .001$). Diabetes: HbA _{1c} levels mean difference = -0.36 ($p = .016$).	Depression: $\Delta 0.610$ Diabetes: $\Delta 0.168$	Significantly greater improvement in depressive symptoms and glycemic control in boxing intervention group versus control group.	Sub	Yes	No
Depression: PHQ9 scores mean difference = -2.14, ($p = .03$) Diabetes: HbA _{1c} levels mean difference = -0.06% ($p = .83$) n.s.	Depression: $\Delta 0.342$ Diabetes: $\Delta -0.032$	Significantly greater improvement in depressive symptoms but not glycemic control in HOPE telehealth intervention versus care as usual control group.	MDD	No	No
Depression: IDS scores mean difference = -3.9 ($p = .248$) n.s. Diabetes: HbA _{1c} levels mean difference = 1.9 ($p = .116$) n.s.	Depression: $\Delta 0.722$ Diabetes: $\Delta -0.032$	Light therapy was not significantly better at reducing depressive symptoms in comparison to placebo, and had no effect on glycemic control.	MDD	No	No

Higgins et al., 2019; Johnson et al., 2014; Nefs et al., 2012; Pibernik-Okanović et al., 2011; Pibernik-Okanovic et al., 2008; Rodríguez et al., 2012; Simson et al., 2008) with low risk of bias however showed a similar effect size: 0.402 (95% CI 0.271;0.533), $p < .0001$ on the combined measures outcome, compared with 0.485 in the original analysis. I^2 was 47, which shows that focusing on low risk-of-bias studies provides similar results but reduces heterogeneity levels (Appendix pp. 20). A Begg funnel plot test for publication bias

with observed and imputed studies showed no small study effect (Appendix pp 21).

6 | DISCUSSION

This systematic review and meta-analysis shows beneficial treatment effects for comorbid depression in type 1 and type 2 diabetes




FIGURE 2 Forest plot showing results of meta-analysis of studies of depressive disorder with depression as outcome, grouped by treatment. CBT = cognitive behavioral therapy; BA = behavioral activation; EX = add-on exercise; PT = psychological treatment (counseling); SPT = supportive psychotherapy

with a larger effect size (0.485) than in the original study that reported an effect size of 0.370 (95% CI 0.470; 0.271) (Van der Feltz-Cornelis et al., 2010). This can be explained by the introduction of new interventions over the last decade with large effect sizes, such as group-based therapy and online treatment. With the emergence of technological developments and increase in accessibility to the internet, treatments delivered online or using mobile technologies have increased in recent years. For example, many psychotherapies such as CBT can now be delivered online. This is particularly useful for people with diabetes given the propensity for poor health outcomes and high healthcare costs in this population.

Also, the effect size of collaborative care increased from a small to moderate effect size as this treatment model has developed over the last decade, especially in the domain of comorbid long-term physical conditions and comorbid depression (O'Hagan & Boreham, 2013; Panagioti et al., 2016; Tully & Baumeister, 2015). There are large differences between treatment effects for different interventions in terms of diabetes and depression outcomes. All interventions

improved depression outcomes significantly in depressive disorder with large effect sizes in group-based therapy, online treatment, exercise, pharmacological treatment and psychotherapy, and moderate effect sizes in collaborative care and phone treatment. However, the effect sizes of such treatments for glycemic control were large in case of pharmacological treatment, group-based therapy and psychotherapy, smaller for collaborative care, and not effective at all in case of exercise, online treatment and phone treatment. The finding that exercise was effective in terms of depression outcomes, but ineffective in improving glycemic control counters expectations for this intervention, as exercise is recommended as a treatment of both type 1 and type 2 diabetes. All current guidelines for depression and diabetes recommend exercise and other aspects of health lifestyle as a first step; this review and meta-analysis, however, shows that exercise is only effective in improving depression. Exercise has been found to be an effective treatment for type 2 diabetes, helping to stabilize plasma glucose and improve body composition, insulin resistance, and glycated hemoglobin. Engagement in exercise is, however,




FIGURE 3 Forest plot showing results of meta-analysis of studies of depressive disorder on glycemic control grouped by treatment. CBT = cognitive behavioral therapy; BA = behavioral activation; EX = add-on exercise PT = psychological treatment (counseling)




FIGURE 4 Flowchart showing treatment recommendations for comorbid depression in diabetes

suboptimal in people with diabetes (Koopmans et al., 2009), and this may be worse in case of comorbid depression (Katon et al., 2010; Lysy et al., 2008). As the findings in this meta-analysis were only based on one study (Groot et al., 2019) on exercise, further research is needed. It would be of interest to assess what the additional effect of an exercise intervention embedded into treatment for diabetes and depression may be. This should be explored in further research as well as the effect of exercise as stand-alone intervention.

This review also shows that interventions that are effective in depressive disorder may not be as effective in subthreshold depression. In this group, psychotherapy and online treatment had large, significant effect sizes on depressive symptoms, but group therapy and psychoeducation were not effective. Looking at glycemic control as an outcome, psychotherapy had a large, significant effect and group-based therapy had a small, significant effect, while online treatment and psychoeducation had no significant effect at all. Consequently, the preferred treatment for both depression and glycemic control in comorbid subthreshold depression would be psychotherapy.

The finding that psychoeducation is not more effective than CAU in subthreshold depression, both for depression outcome and glycemic control, is an important finding as in stepped care models, psychoeducation has been suggested as a first step in diabetes-related distress or subthreshold depressive symptoms (Huang et al., 2013). Furthermore, psychoeducation was supposed to be a good start for improving self-management and in that way improving glycemic control. This line of thought is not supported by our results. Also, the finding that group therapy is highly effective in depressive disorder, but not in subthreshold depression, might suggest that patients with subthreshold depression might benefit more from individual treatment tailored to their specific needs rather than from group participation, something that has been suggested earlier (Huang et al., 2013). Treatment of comorbid subthreshold depressive disorder could be psychotherapy both in patients with elevated or normal HbA_{1c}. The latter group might also benefit from online treatment. If glycemic control is a target, our analysis shows that it makes sense to target patients with high baseline levels of depression and of HbA_{1c}, as they are likely to benefit most from treatment on both symptom levels.

In our flowchart, we recommend collaborative care in comorbid MDD and multimorbidity or problems requiring complex case management. Although effect sizes for some other treatment modes are found to be larger in our meta-analysis, none of those were evaluated in patients with such a complex and multimorbid profile, whereas several systematic reviews show that outcomes in such patient groups improve by collaborative care (Faridhosseini et al., 2014; Tully & Baumeister, 2015).

One RCT (Guo et al., 2014) found that metformin improved glycemic control but also depressive outcomes, compared to placebo, in patients with type 2 diabetes. Although a small study with only 58 participants, this finding is of interest and may contribute to the expanding field of evaluation of medicines that are normally prescribed for physical conditions for their effect in treatment of depression

(Arteaga-Henríquez et al., 2019; Che et al., 2018; Köhler et al., 2014). Further research could explore the mechanism for metformin in improvement of depression in diabetes.

Our study has several strengths. First, we included data without language restriction from studies identified by a comprehensive search of the published literature. We included studies exploring the effect of treatment in subthreshold depression. Our sensitivity analysis excluding high risk-of-bias studies confirmed the findings, the fixed model meta-analysis refuted the null hypothesis, and we found no indication for publication bias. Second, we provided relative effect sizes for several new treatment modalities compared to the treatments already explored in the first systematic review, we differentiated the treatment effect on depressive outcomes versus glycemic control, and by performing meta-regression we showed the influence of baseline depression severity on both depression outcome and glycemic control, whereas baseline HbA_{1c} only influenced glycemic control as an outcome. This combination of findings enabled us to provide clinicians with innovative guidance about which interventions may suit best, depending on patient profile. These strengths make our study the most comprehensive systematic review and meta-analysis of treatment for comorbid depression in diabetes yet undertaken.

Our analysis has several limitations. First, most of the included studies did not meet all criteria to reduce risk of bias, mostly due to unclear reporting and to small samples. Despite our efforts to contact authors for missing data, we were unable to include such data in three studies due to lack of response (Eli et al., 2011; Petrak, Herpertz, et al., 2015; van der Sluijs et al., 2018), which may have to do with the long timeframe of this systematic review. The need for low risk-of-bias studies in this field remains, with proper reporting of methodology and of outcomes. Second, the planned moderator analyses on the effect of add-on exercise on treatment outcome and on adherence as an outcome of treatment could not be performed because of insufficient data (Appendix pp. 22). Third, some treatments were only evaluated in one RCT. This probably reflects that, although many of these “new treatments” have been used for some time and have been felt to be useful by patients and clinicians, at least in primary care, researchers had not actively examined these “new” treatments until recently. In view of their clinical relevance, we emphasize this limitation. We strongly suggest further research is needed especially in group-based treatment and exercise, that seem to have promising results. Another limitation concerns the provenance of the studies. Although this is a study with a global perspective in terms of included studies, it is clear that there is a scarcity of data from many low- and middle-income countries, as shown by the map in Figure 1. The imbalance is of growing importance because it is likely that the low- and middle-income countries will have the greatest increases of comorbidities of prevalence and incidence of diabetes and depression. In countries in which the attention to mental health problems is minimal or absent and the investment in the care for diabetes is appropriate, the guidance for treatment that we could deduce from this systematic review and meta-analysis is particularly relevant and may improve care for comorbid depression.

Furthermore, the studies in this meta-analysis do not present results for type 1 diabetes and type 2 diabetes separately despite the different types of diabetes affecting different groups of the population; for example, type 2 diabetes tends to develop more commonly in older people compared with a peak incidence of type 1 diabetes in adolescence and young adulthood. The lack of studies in type 1 diabetes alone with comorbid depression or comorbid subthreshold disorder is striking and research is needed to fill this gap.

A clearer understanding of the mechanisms underpinning why some treatments are more effective for patients with depressive disorder than for subthreshold depression and vice versa would also greatly benefit this area of research and for this purpose studies might provide more detailed information about the contents of the intervention. In particular, the idea that interventions aiming to improve self-management lead to better adherence and better diabetes and depression outcomes should be challenged in research as studies reporting on adherence as an outcome are lacking. Studies are also needed to develop standardized techniques or tools to help better identify particular subtypes of patients taking into account their depression severity and glycemic control. These suggestions will further aid in the identification and personalization of appropriate treatment plans for patients with diabetes and depression as outlined above.

ACKNOWLEDGMENT

Not Applicable.

CONFLICTS OF INTEREST

Over the last three years, RIGH has received honoraria for speaker engagement, conference attendance, or advisory boards from AstraZeneca, Boehringer Ingelheim, European Association for the Study of Diabetes, Eli Lilly, Janssen, Menarini, Mylan, Novo Nordisk, OmniaMed, and Otsuka. AN has received funding for lectures from OmniaMed, and, as Chairperson of the Psycho-Social Aspects of Diabetes (PSAD) study group of the European Association for the Study of Diabetes, funding for travel Fellowships for early career researchers from Sanofi. The other authors have no conflicts of interest to declare.

AUTHOR CONTRIBUTIONS

This systematic review was designed by CFC and co-authors. Screening and data extraction were completed by SA and CFC. The meta-analysis was performed by CFC. The initial version of the manuscript was written by SA and CFC. Following this, input was provided from all authors. All authors approved the final version of the manuscript.

FUNDING INFORMATION

This study was financially supported by Hull York Medical School. The funder had no role in study design, data collection, data analysis, data interpretation, writing of the report, and decision to submit the paper for publication. The corresponding author had full access to all data in the study and had final responsibility for the decision to submit for publication.

ETHICAL APPROVAL

Ethical approval was not required for the current study as the data entered in the meta-analysis were collated from previous clinical trials in which informed consent had already been obtained.

DATA AVAILABILITY STATEMENT

Data available on request.

ORCID

Christina van der Feltz-Cornelis  <https://orcid.org/0000-0001-6925-8956>

[org/0000-0001-6925-8956](https://orcid.org/0000-0001-6925-8956)

Sarah F. Allen  <https://orcid.org/0000-0003-0274-4137>

Richard I. G. Holt  <https://orcid.org/0000-0001-8911-6744>

Arie Nouwen  <https://orcid.org/0000-0002-0609-4082>

Norman Sartorius  <https://orcid.org/0000-0001-8708-6289>

REFERENCES

- Arteaga-Henríquez, G., Simon, M. S., Burger, B., Weidinger, E., Wijkhuijs, A., Arolt, V., Birkenhager, T. K., Musli, R., Müller, N., & Drexhage, H. A. (2019). Low-grade inflammation as a predictor of antidepressant and anti-inflammatory therapy response in MDD patients: A systematic review of the literature in combination with an analysis of experimental data collected in the EU-Moodinflamm consortium. *Frontiers Psychiatry, 10*. <https://doi.org/10.3389/fpsy.2019.00458>
- Atlantis, E., Fahey, P., & Foster, J. (2014). Collaborative care for comorbid depression and diabetes: A systematic review and meta-analysis. *British Medical Journal Open, 4*(4), e004706.
- Barragán-Rodríguez, L., Rodríguez-Morán, M., & Guerrero-Romero, F. (2008). Efficacy and safety of oral magnesium supplementation in the treatment of depression in the elderly with type 2 diabetes: A randomized, equivalent trial. *Magnesium Research, 21*(4), 218–223.
- Baumeister, H. H. N., & Bengel, J. (2012). Psychological and pharmacological interventions for depression in patients with diabetes mellitus and depression. *Cochrane Database of Systematic Reviews (Online), 12*, CD008381.
- Baumeister, H., Hutter, N., & Bengel, J. (2014). Psychological and pharmacological interventions for depression in patients with diabetes mellitus: An abridged Cochrane review. *Diabetic Medicine, 31*(7), 773–786.
- Begg, C. B. (1994). Publication bias. *The Handbook of Research Synthesis, 25*, 299–409.
- Beydoun, M. A., & Wang, Y. (2010). Pathways linking socioeconomic status to obesity through depression and lifestyle factors among young US adults. *Journal of Affective Disorders, 123*, 52–63.
- Biostat (2005). *Comprehensive Meta-analysis, Version 2. [computer program]*. : Biostat.
- Black, S. A., Markides, K. S., & Ray, L. A. (2003). Depression predicts increased incidence of adverse health outcomes in older Mexican Americans with type 2 diabetes. *Diabetes Care, 26*(10), 2822–2828.
- Bogner, H. R., Morales, K. H., de Vries, H. F., & Cappola, A. R. (2012). Integrated management of type 2 diabetes mellitus and depression treatment to improve medication adherence: A randomized controlled trial. *Annals of Family Medicine, 10*(1), 15–22. <https://doi.org/10.1370/afm.1344>
- Bot, M. P. F., Ormel, J., Slaets, J. P., & De Jonge, P. (2010). Predictors of incident major depression in diabetic outpatients with subthreshold depression. *Diabetic Medicine, 27*, 1295–1301.
- Brouwer, A., van Raalte, D. H., Nguyen, H. T., Rutters, F., van de Ven, P., Elders, P. J. M., Moll, A. C., Van Someren, E. J. W., Snoek, F. J., Beekman, A. T. F., & Brenner, M. A. (2019). Effects of light therapy on mood and insulin sensitivity in patients with type 2 diabetes

- and depression: results from a randomized placebo-controlled trial. *Diabetes Care*, 42(4), 529–538. <https://doi.org/10.2337/dc18-1732>
- Champaneri, S. W. G., Malhotra, S. S., Casagrande, S. S., & Golden, S. H. (2010). Biological basis of depression in adults with diabetes. *Current Diabetes Reports*, 10, 396–405.
- Che, T., Teng, X., Huang, Q., Mu, Y., Tang, X., Mu, X., & Wei, Y. (2018). Agomelatine versus fluoxetine in glycemic control and treating depressive and anxiety symptoms in type 2 diabetes mellitus subjects: A single-blind randomized controlled trial. *Neuropsychiatric Disease and Treatment*, 14, 1527.
- Ciechanowski, P. S., Katon, W. J., & Russo, J. E. (2000). Depression and diabetes: Impact of depressive symptoms on adherence, function, and costs. *Archives of Internal Medicine*, 160(21), 3278–3285. <https://doi.org/10.1001/archinte.160.21.3278>
- de Groot, M., Anderson, R., Freedland, K. E., Clouse, R. E., & Lustman, P. J. (2001). Association of depression and diabetes complications: A meta-analysis. *Psychosomatic Medicine*, 63(4), 619–630. <https://doi.org/10.1097/00006842-200107000-00015>
- de Groot, M., Shubrook, J. H., Hornsby, W. G. Jr, Pillay, Y., Mather, K. J., Fitzpatrick, K., Yang, Z., & Saha, C. (2019). Program ACTIVE II: outcomes from a randomized, multistate community-based depression treatment for rural and Urban adults With Type 2 diabetes. *Diabetes Care*, 42(7), 1185–1193. <https://doi.org/10.2337/dc18-2400>
- DerSimonian, R., & Laird, N. (1986). Meta-analysis in clinical trials. *Controlled Clinical Trials*, 7(3), 177–188. [https://doi.org/10.1016/0197-2456\(86\)90046-2](https://doi.org/10.1016/0197-2456(86)90046-2)
- Ebert, D. D., Nobis, S., Lehr, D., Bauemeister, H., Ripe, H., Auerbach, R. P., Snoek, F., Cuijpers, P., & Berking, M. (2017). The 6-month effectiveness of Internet-based guided self-help for depression in adults with Type 1 and 2 diabetes mellitus. *Diabetic Medicine*, 34(1), 99–107. <https://doi.org/10.1111/dme.13173>
- Echeverry, D., Duran, P., Bonds, C., Lee, M., & Davidson, M. B. (2009). Effect of pharmacological treatment of depression on A1C and quality of life in low-income Hispanics and African Americans with diabetes: A randomized, double-blind, placebo-controlled trial. *Diabetes Care*, 32(12), 2156–2160. <https://doi.org/10.2337/dc09-0785>
- Ell, K., Katon, W., Xie, B., Lee, P. J., Kapetanovic, S., Guterman, J., & Chou, C. P. (2010). Collaborative care management of major depression among low-income, predominantly Hispanic subjects with diabetes: A randomized controlled trial. *Diabetes Care*, 33(4), 706–713. <https://doi.org/10.2337/dc09-1711>
- Ell, K., Katon, W., Xie, B., Lee, P. J., Kapetanovic, S., Guterman, J., & Chou, C. P. (2011). One-year postcollaborative depression care trial outcomes among predominantly Hispanic diabetes safety net patients. *General Hospital Psychiatry*, 33(5), 436–442. <https://doi.org/10.1016/j.genhosppsy.2011.05.018>
- Faridhosseini, F., Sadeghi, R., Farid, L., & Pourgholami, M. (2014). Celecoxib: A new augmentation strategy for depressive mood episodes. A systematic review and meta-analysis of randomized placebo-controlled trials. *Human Psychopharmacology: Clinical and Experimental*, 29(3), 216–223.
- GBD 2017 Disease and Injury Incidence and Prevalence Collaborators (2018). Global Burden of Disease 2016 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990–2017: A systematic analysis for the Global Burden of Disease Study 2017. *The Lancet* 392(10100), 1211–1259.
- Gois, C., Dias, V., Carmo, I., Duarte, R., Ferro, A., Santos, A. L., Sousa, F., & Barbosa, A. (2014). Treatment response in type 2 diabetes patients with major depression. *Clinical Psychology & Psychotherapy*, 21(1), 39–48. <https://doi.org/10.1002/cpp.1817>
- Golden, S. H., Lazo, M., Carnethon, M., Bertoni, A. G., Schreiner, P. J., Roux, A. V. D., Lee, H. B., & Lyketsos, C. (2008). Examining a bidirectional association between depressive symptoms and diabetes. *JAMA*, 299(23), 2751–2759.
- Gonzalez, J. S., Peyrot, M., McCarl, L. A., Collins, E. M., Serpa, L., Mimiaga, M. J., & Safren, S. A. (2008). Depression and diabetes treatment nonadherence: A meta-analysis. *Diabetes Care*, 31(12), 2398–2403. <https://doi.org/10.2337/dc08-1341>
- Gonzalez, J. S., Safren, S., Delahanty, L., Cagliero, E., Wexler, D. J., Meigs, J. B., & Grant, R. W. (2008). Symptoms of depression prospectively predict poorer self-care in patients with Type 2 diabetes. *Diabetic Medicine*, 25(9), 1102–1107.
- Gulseren, L., Gulseren, S., Hekimsoy, Z., & Mete, L. (2005). Comparison of fluoxetine and paroxetine in type II diabetes mellitus patients. *Archives of Medical Research*, 36(2), 159–165. <https://doi.org/10.1016/j.arcmed.2004.12.018>
- Guo, M., Mi, J., Jiang, Q. M., Xu, J. M., Tang, Y. Y., Tian, G., & Wang, B. (2014). Metformin may produce antidepressant effects through improvement of cognitive function among depressed patients with diabetes mellitus. *Clinical and Experimental Pharmacology and Physiology*, 41(9), 650–656. <https://doi.org/10.1111/1440-1681.12265>
- Guyatt, G. H., Oxman, A., Vist, G. E., Kunz, R., Falck-Ytter, Y., Alonso-Coello, P., & Schunemann, H. J. (2008). GRADE: An emerging consensus on rating quality of evidence and strength of recommendations. *BMJ (Clinical Research ed)*, 336(7650), 924–926. <https://doi.org/10.1136/bmj.39489.470347.AD>
- Harkness, E. M. W., Valderas, J., Coventry, P., Gask, L., & Bower, P. (2010). Identifying psychosocial interventions that improve both physical and mental health in patients with diabetes: A systematic review and meta-analysis. *Diabetes Care*, 33, 926–930.
- Hermanns, N., Schmitt, A., Gahr, A., Herder, C., Nowotny, B., Roden, M., Ohmann, C., Kruse, J., Haak, T., & Kulzer, T. (2015). The effect of a diabetes-specific cognitive behavioral treatment program (DIAMOS) for patients with diabetes and subclinical depression: Results of a randomized controlled trial. *Diabetes Care*, 38(4), 551–560.
- Higgins, J. P. T. (2003). Measuring inconsistency in meta-analyses. *BMJ*, 327(7414), 557–560
- Higgins, J. P. T. J., Chandler, J., Cumpston, M., Li, T., Page, M. J., Welch, V. A. (2019). *Cochrane handbook for systematic reviews of interventions*, 2nd edn. : John Wiley and Sons.
- Hofmann, M. K. B., Leichsenring, F., & Kruse, J. (2013). Depression as a risk factor for mortality in individuals with diabetes: A meta-analysis of prospective studies. *PLoS One*, 8(11), e79809.
- Huang, C. Y., Lai, H. L., Chen, C. I., Lu, Y. C., Li, S. C., Wang, L. W., & Su, Y. (2016). Effects of motivational enhancement therapy plus cognitive behaviour therapy on depressive symptoms and health-related quality of life in adults with type II diabetes mellitus: A randomised controlled trial. *Quality of Life Research*, 25(5), 1275–1283.
- Huang, X., Song, L., Li, T., Li, J., Wu, S., & Li, N. (2002). Effect of health education and psychosocial intervention on depression in patients with Type II diabetes. *Chinese Mental Health Journal*, 16, 149–151.
- Huang, Y. W. X., Wu, T., Chen, R., & Guo, A. (2013). Collaborative care for patients with depression and diabetes mellitus: A systematic review and meta-analysis. *BMC Psychiatry* 13, 260.
- Johnson, J. A., Al Sayah, F., Wozniak, L., Rees, S., Soprovich, A., Qiu, W., Chik, C. L., Chue, P., Florence, P., Jacquier, J., Lysak, P., Opgenorth, A., Katon, W., & Majumdar, S. R. (2014). Collaborative care versus screening and follow-up for patients with diabetes and depressive symptoms: results of a primary care-based comparative effectiveness trial. *Diabetes Care*, 37(12), 3220–3226.
- Joseph, J. J., & Golden, S. H. (2017). Cortisol dysregulation: The bidirectional link between stress, depression, and type 2 diabetes mellitus. *Annals of the New York Academy of Sciences*, 1391(20), 1391.
- Kang, R., He, Y., Yan, Y., Li, Z., Wu, Y., Guo, X., Liang, Z., & Jiang, J. (2015). Comparison of paroxetine and agomelatine in depressed type 2 diabetes mellitus patients: A double-blind, randomized, clinical trial. *Neuropsychiatric Disease and Treatment*, 11, 1307–1311.

- Karaiskos, D., Tzavellas, E., Ilias, I., Liappas, I., & Paparrigopoulos, T. (2013). Agomelatine and metformin for the treatment of depression in type 2 diabetes mellitus. *International Journal of Clinical Practice*, 67(3), 257–260.
- Katon, W. J. (2008). The comorbidity of diabetes mellitus and depression. *American Journal of Medicine*, 121(11 Suppl 2), S8–15.
- Katon, W., Lin, E. H., & Kroenke, K. (2007). The association of depression and anxiety with medical symptom burden in patients with chronic medical illness. *General Hospital Psychiatry*, 29(2), 147–155.
- Katon, W., & Van der Feltz-Cornelis, C. M. (2010). Treatment of depression in patients with diabetes: Efficacy, effectiveness and maintenance trials, and new service models. In W. Katon, N. Maj, & N. Sartorius (Eds.), *Depression and diabetes* (pp. 81–108). Wiley.
- Katon, W. J., Von Korff, M., Lin, E. H., Simon, G., Ludman, E., Russo, J., Ciechanowski, P., Walker, E., & Bush, T. (2004). The Pathways Study: A randomized trial of collaborative care in patients with diabetes and depression. *Archives of General Psychiatry*, 61(10), 1042–1049.
- Katon, W. J., Von Korff, M., Lin, E. H., Simon, G., Ludman, E., Russo, J., Ciechanowski, P., Walker, E., & Bush, T. (2004). The pathways study: A randomized trial of collaborative care in patients with diabetes and depression. *Archives of General Psychiatry*, 61(10), 1042–1049. <https://doi.org/10.1001/archpsyc.61.10.1042>
- Khaledi, M., Haghghatdoost, F., Feizi, A., & Aminorroaya, A. (2019). The prevalence of comorbid depression in patients with type 2 diabetes: An updated systematic review and meta-analysis on huge number of observational studies. *Acta Diabetologica*, 56(6), 631–650. <https://doi.org/10.1007/s00592-019-01295-9>
- Khazaie, H., Rahimi, M., Tatari, F., Rezaei, M., Najafi, F., & Tahmasian, M. (2011). Treatment of depression in type 2 diabetes with fluoxetine or citalopram. *Neurosciences (Riyadh)*, 16(1), 42–45.
- Geraets, A. F. J., Köhler, S., Muzambi, R., Schalkwijk, C. G., Oenema, A., Eussen, S. J. P. M., Dagnelie, P. C., Stehouwer, C. D. A., Schaper, N. C., Henry, R. M. A., van der Kallen, C. J. A., Wesselius, A., Koster, A., Verhey, F., & Schram, M. T. (2020). The association of hyperglycaemia and insulin resistance with incident depressive symptoms over 4 years of follow-up: The Maastricht Study. *Diabetologia*, 63, 2315–2328.
- Köhler, O., Benros, M. E., Nordentoft, M., Farkouh, M. E., Lyengar, R. L., Mors, O., & Krogh, J. (2014). Effect of anti-inflammatory treatment on depression, depressive symptoms, and adverse effects: A systematic review and meta-analysis of randomized clinical trials. *JAMA Psychiatry*, 71(12), 1381–1391.
- Koopmans, B. P. F., de Bie, R. A., van Rooij, E. S., Leusink, G. L., & Pop, V. J. (2009). Depressive symptoms are associated with physical inactivity in patients with type 2 diabetes. The DIAZOB Primary Care Diabetes study. *Family Practice*, 26, 171–173.
- Kumar, K., Salman, M., Shukla, V., Ahmad, A., Kumar Verma, V., Pizvi, D., & Kohli, A. (2015). Comparative effect of agomelatine versus escitalopram on glycemic control and symptoms of depression in patients with type 2 diabetes mellitus and depression. *JPSR*, 6(10), 4304–4309.
- Laake, J.-P.-S., Stahl, D., Amiel, S. A., Petrak, F., Sherwood, R. A., Pickup, J. C., & Ismail, K. (2014). The association between depressive symptoms and systemic inflammation in people with type 2 diabetes: Findings from the South London Diabetes Study. *Diabetes Care*, 37, 2186–2192.
- Li, S. C., Li, M., & Song, S. J. (2003). The effect of psychological intervention in treating the diabetic patients with negative emotion. *Shandong Journal of Psychological Medicine*, 16, 148.
- Liberati, A., Altman, D. G., Tetzlaff, J., Mulrow, C., Gotzsche, P. C., Ioannidis, J. P., Clarke, M., Devereaux, P. J., Kleijnen, J., & Moher, D. (2009). The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: Explanation and elaboration. *PLoS Med*, 6(7), e1000100. <https://doi.org/10.1371/journal.pmed.1000100>
- Lipsey, M. W., & Wilson, D. B. (1993). The efficacy of psychological, educational, and behavioral treatment: Confirmation from meta-analysis. *American Psychologist*, 48(12), 1181. <https://doi.org/10.1037/0003-066X.48.12.1181>
- Lloyd, C. E., Nouwen, A., Sartorius, N., Ahmed, H. U., Alvarez, A., Bahendeka, S., Basangwa, D., Bobrov, A. E., Boden, S., Bulgari, V., Burti, L., Chaturvedi, S. K., Cimino, L. C., Gaebel, W., de Girolamo, G., Gondek, T. M., de Braude, M. G., Guntupalli, A., Heinze, M. G., ... Xin, Y. (2018). Prevalence and correlates of depressive disorders in people with Type 2 diabetes: Results from the International Prevalence and Treatment of Diabetes and Depression (INTERPRET-DD) study, a collaborative study carried out in 14 countries. *Diabetic Medicine* 35(6), 760–769.
- Long, F., Yan, J., Hu, P., Xia, M., Liu, H., & Gu, C. (2015). Effect of group counseling on depression, compliance and blood sugar level in diabetic patients. *Journal of Central South University Medical Sciences*, 40(8), 879–885.
- Lu, X., Lu, B., & Gu, X. (2005). Cognitive therapy in combination with electromyographic feedback in treatment of diabetes patients with depression after cerebral infarction. *Chinese Journal of Clinical Psychology*, 14, 215–216.
- Lustman, P. J., Anderson, R. J., Freedland, K. E., de Groot, M., Carney, R. M., & Clouse, R. E. (2000). Depression and poor glycemic control: A meta-analytic review of the literature. *Diabetes Care*, 23(7), 934–942. <https://doi.org/10.2337/diacare.23.7.934>
- Lustman, P. J., Freedland, K. E., Griffith, L. S., & Clouse, R. E. (2000). Fluoxetine for depression in diabetes: A randomized double-blind placebo-controlled trial. *Diabetes Care*, 23(5), 618–623.
- Lustman, P. J., Griffith, L. S., Clouse, R. E., Freedland, K. E., Eisen, S. A., Rubin, E. H., Carney, R. M., & McGill, J. B. (1997). Effects of nortriptyline on depression and glycemic control in diabetes: Results of a double-blind, placebo-controlled trial. *Psychosomatic Medicine*, 59(3), 241–250.
- Lustman, P. J., Griffith, L. S., Freedland, K. E., Kissel, S. S., & Clouse, R. E. (1998). Cognitive behavior therapy for depression in type 2 diabetes mellitus. A randomized, controlled trial. *Annals of Internal Medicine*, 129(8), 613–621.
- Lustman, P. J., Griffith, L. S., Freedland, K. E., Kissel, S. S., & Clouse, R. E. (1998). Cognitive behavior therapy for depression in type 2 diabetes mellitus: A randomized, controlled trial. *Annals of Internal Medicine*, 129(8), 613–621.
- Lustman, P. J., Williams, M. M., Sayuk, G. S., Nix, B. D., & Clouse, R. E. (2007). Factors influencing glycemic control in type 2 diabetes during acute- and maintenance-phase treatment of major depressive disorder with bupropion. *Diabetes Care*, 30(3), 459–466.
- Lysy, Z., Da, C. D., & Dasgupta, K. (2008). The association of physical activity and depression in Type 2 diabetes. *Diabetic Medicine*, 25(10), 1133–1141.
- McGuire, H., Moncrieff, J., & Churchill, R. (1998). *Quality assessment of trials found within the scope of Cochrane Collaboration Depression, Anxiety and Neurosis (CCDAN)*. International Cochrane Colloquium.
- Naik, A. D., Hundt, N. E., Vaughan, E. M., Petersen, N. J., Zeno, D., Kunik, M., & Cully, J. A. (2019). Effect of telephone-delivered collaborative goal setting and behavioral activation vs enhanced usual care for depression among adults with uncontrolled diabetes: a randomized clinical trial. *JAMA Network Open*, 2(8), e198634.
- Nefs, G., Pouwer, F., Denollet, J., & Pop, V. (2012). The course of depressive symptoms in primary care patients with type 2 diabetes: Results from the Diabetes, Depression, Type D Personality Zuidoost-Brabant (DiaDDZoB) Study. *Diabetologia*, 55, 608–616.
- Newby, J., Robins, L., Wilhelm, K., Smith, J., Fletcher, T., Gillis, I., Ma, T., Finch, A., Campbell, L., & Andrews, G. (2017). Web-based cognitive behavior therapy for depression in people with diabetes mellitus: a randomized controlled trial. *Journal of Medical Internet Research*, 19(5), e157.

- Nouwen, A., Adriaanse, M. C., van Dam, K., Iversen, M. M., Viechtbauer, W., Peyrot, M., Caramlau, I., Kokoszka, A., Kanc, K., de Groot, M., Nefs, G., & Pouwer, F. (2019). European Depression in Diabetes (EDID) Research Consortium. Longitudinal associations between depression and diabetes complications: A systematic review and meta-analysis. *Diabetic Medicine*, *36*, 1562–1572.
- O'Hagan, C. D. V. G., & Boreham, C. A. (2013). Exercise prescription in the treatment of type 2 diabetes mellitus. *Sports Medicine*, *43*, 39–49.
- O'Connor, P. J., Crain, A. L., Rush, W. A., Hanson, A. M., Fischer, L. R., & Kluznik, J. C. (2009). Does diabetes double the risk of depression? *The Annals of Family Medicine*, *7*(4), 328–335.
- Paile-Hyvärinen, M., Wahlbeck, K., & Eriksson, J. G. (2003). Quality of life and metabolic status in mildly depressed women with type 2 diabetes treated with paroxetine: A single-blind randomised placebo controlled trial. *BMC Family Practice*, *4*(1), 7.
- Paile-Hyvärinen, M., Wahlbeck, K., & Eriksson, J. G. (2007). Quality of life and metabolic status in mildly depressed patients with type 2 diabetes treated with paroxetine: A double-blind randomised placebo controlled 6-month trial. *BMC Family Practice*, *8*, 34. <https://doi.org/10.1186/1471-2296-8-34>
- Panagiotti, M., Bower, P., Lovell, K. E., & Gilbody, S., Waheed, W., Dickens, C., Archer, J., Simon, G., Ell, K., Huffman, J. C., Richards, D. A., van der Feltz-Cornelis, C., Adler, D. A., Bruce, M., Buszewicz, M., Cole, M. G., Davidson, K. W., de Jonge, P., Gensichen, J., ... Coventry, P. A. (2016). Association between chronic physical conditions and the effectiveness of collaborative care for depression: An individual participant data meta-analysis. *JAMA. Psychiatry*, *73*(9), 978–989.
- Penckofer, S. M., Ferrans, C., Mumby, P., Byrn, M., Emanuele, M. A., Harrison, P. R., & Durazo-Arvizu, R. A., & Lustman, P. (2012). A psychoeducational intervention (SWEEP) for depressed women with diabetes. *Annals of Behavioral Medicine*, *44*(2), 192–206.
- Petrak, F., Baumeister, H., Skinner, T. C., Brown, A., & Holt, R. I. G. (2015). Depression and diabetes: Treatment and health-care delivery. *Review Lancet Diabetes Endocrinol* *6*, 472–485.
- Petrak, F., Herpertz, S., Albus, C., Hermanns, N., Hiemke, C., Hiller, W., Kronfeld, K., Kruse, J., Kulzer, B., Ruckes, C., Zahn, D., & Müller, M. J. (2015). Cognitive behavioral therapy versus sertraline in patients with depression and poorly controlled diabetes: The diabetes and depression (DAD) study: A randomized controlled multicenter trial. *Diabetes Care*, *38*(5), 767–775. <https://doi.org/10.2337/dc14-1599>
- Pibernik-Okanović, M., Ajduković, D., Lovrenčić, M. V., & Hermanns, N. (2011). Does treatment of subsyndromal depression improve depression and diabetes related outcomes: Protocol for a randomised controlled comparison of psycho-education, physical exercise and treatment as usual. *Trials*, *12*(1), 17.
- Pibernik-Okanovic, M., Begic, D., Ajdukovic, D., Andrijasevic, N., & Metelko, Z. (2009). Psychoeducation versus treatment as usual in diabetic patients with subthreshold depression: Preliminary results of a randomized controlled trial. *Trials*, *10*, 78.
- Pibernik-Okanovic, M., Begic, D., Peros, K., Szabo, S., & Metelko, Z. (2008). Psychosocial factors contributing to persistent depressive symptoms in type 2 diabetic patients: A Croatian survey from the European Depression in Diabetes Research Consortium. *Journal of Diabetes and Its Complications*, *22*(4), 246–253.
- Pibernik-Okanovic, M., Hermanns, N., Ajdukovic, D., Kos, J., Prasek, M., Sekerija, M., & Vucuc, L. M. (2015). Does treatment of subsyndromal depression improve depression-related and diabetes-related outcomes? A randomised controlled comparison of psychoeducation, physical exercise and enhanced treatment as usual. *Trials*, *16*, 305.
- Piette, J. D., Richardson, C., Himle, J., Duffy, S., Torres, T., Vogel, M., Barber, K., & Valenstein, K. (2011). A randomized trial of telephone counseling plus walking for depressed diabetes patients. *Medical Care* *49*(7), 641.
- Renn, B. N., Feliciano, L., & Segal, D. L. (2011). The bidirectional relationship of depression and diabetes: A systematic review. *Clinical Psychology Review*, *31*, 1239–1246. <https://doi.org/10.1016/j.cpr.2011.08.001>
- Rodríguez, M. R., Nuevo, R., Chatterji, S., & Ayuso-Mateos, J. L. (2012). Definitions and factors associated with subthreshold depressive conditions: A systematic review. *BMC Psychiatry*, *12*(1), 181.
- Rothstein, H. R., Sutton, A. J., & Borenstein, M. (Eds.) (2005). Publication bias in meta-analysis: Prevention, assessment and adjustments. In *Publication bias in meta-analysis* (pp. 1–7). John Wiley & Sons.
- Ruengorn, C. S. K., Niwatananun, W., Mahatnirunkul, S., Pumpaisalchai, W., & Patumanond, J. (2012). Factors related to suicide attempts among individuals with major depressive disorder. *International Journal of General Medicine*, *5*, 323.
- Safren, S. A., Gonzalez, J. S., Wexler, D. J., Psaros, C., Delahanty, L. M., Blashill, A. J., Margolina, A. I., & Cagliero, E. (2014). A randomized controlled trial of cognitive behavioral therapy for adherence and depression (CBT-AD) in patients with uncontrolled type 2 diabetes. *Diabetes Care*, *37*(3), 625–633.
- Schneider, K. L., Panza, E., Handschin, B., Ma, Y., Busch, A. M., Waring, M. E., Appelhans, B. M., Whited, M. C., Keeney, J., Kern, D., Blendea, M., Ockene, I., & Pagoto, S. L. (2016). Feasibility of pairing behavioral activation with exercise for women with type 2 diabetes and depression: The get it study pilot randomized controlled trial. *Behavior Therapy*, *47*(2), 198–212.
- Simson, U., Nawarotzky, U., Friese, G., Porck, W., Schottenfeld-Naor, Y., Hahn, S., Scherbaum, W. A., & Kruse, J. (2008). Psychotherapy intervention to reduce depressive symptoms in patients with diabetic foot syndrome. *Diabetic Medicine*, *25*(2), 206–212.
- Stoop, C., Nefs, G., Pommer, A., Pop, V., & Pouwer, F. (2015). Effectiveness of a stepped care intervention for anxiety and depression in people with diabetes, asthma or COPD in primary care: A randomized controlled trial. *Journal of Affective Disorders*, *184*, 269–276.
- Tovote, K. A., Fleer, J., Snippe, E., Peeters, A. C. T. M., Emmelkamp, P. M. G., Sanderman, R., Links, T. P., & Schroevers, M. J. (2014). Individual mindfulness-based cognitive therapy (MBCT) and cognitive behavior therapy (CBT) for treating depressive symptoms in patients with diabetes: Results of a randomized controlled trial. *Diabetes Care*, *37*(9), 2427–2434.
- Tully, P. J., & Baumeister, H. (2015). Collaborative care for comorbid depression and coronary heart disease: A systematic review and meta-analysis of randomised controlled trials. *British Medical Journal Open*, *5*(12), e009128.
- van Bastelaar, K. M., Pouwer, F., Cuijpers, P., Riper, H., & Snoek, F. J. (2011). Web-based depression treatment for type 1 and type 2 diabetic patients: A randomized, controlled trial. *Diabetes Care*, *34*(2), 320–325. <https://doi.org/10.2337/dc10-1248>
- Van der Feltz-Cornelis, C. M., & Ader, H. J. (2000). Randomization in psychiatric intervention research in the general practice setting. *International Journal of Methods in Psychiatric Research*, *9*(3), 134–142. <https://doi.org/10.1002/mpr.87>
- Van der Feltz-Cornelis, C. M., Nuyen, J., Stoop, C., Chan, J., Jacobson, A. M., Katon, W., Snoek, F. J., & Sartorius, N. (2010). Effect of interventions for major depressive disorder and significant depressive symptoms in patients with diabetes mellitus: A systematic review and meta-analysis. *General Hospital Psychiatry*, *32*(4), 380–395. <https://doi.org/10.1016/j.genhosppsych.2010.03.011>
- van der Sluijs, J. F. C. H., Eijlsbroek, V., Rijnders, C. A. T., Van Marwijk, H. W. J., & van der Feltz-Cornelis, C. M. (2018). Illness burden and physical outcomes associated with collaborative care in patients with comorbid depressive disorder in chronic medical conditions: A systematic review and meta-analysis. *General Hospital Psychiatry*, *50*, 1–14.
- Van Steenberg-Weijnenburg, K. M. V. P. A., Horn, E. K., Nuyen, J., van Dam, P. S., van Benthem, T. B., Beekman, A. T., Rutten, F. F.,

- Hakkaart-van Roijen, L., & van der Feltz-Cornelis, C. M. (2011). More co-morbid depression in patients with Type-2 diabetes with multiple complications. An observational study at a specialized outpatient clinic. *Diabetic Medicine*, 28(1), 86–89.
- Williams, J. W., Katon, W., Lin, E. H., Noel, P. H., Worchel, J., Cornell, J., Harpole, L., Fultz, B. A., Hunkeler, E., Mika, V. S., & Unutzer, J. (2004). The effectiveness of depression care management on diabetes-related outcomes in older patients. *Annals of Internal Medicine*, 140(12), 1015–1024. <https://doi.org/10.7326/0003-4819-140-12-200406150-00012>
- Xue, H. (2004). Paroxetine for depression in diabetes: A randomized controlled trial. *Chinese Mental Health Journal*, 18, 735–737.
- Zheng, Y., Ley, S. H., & Hu, F. B. (2018). Global aetiology and epidemiology of type 2 diabetes mellitus and its complications. *Nature Reviews Endocrinology*, 14(2), 88. <https://doi.org/10.1038/nrendo.2017.151>
- Zheng, Y., Zhou, Y., & Lai, Q. (2015). Effects of twenty-four move shadow boxing combined with psychosomatic relaxation on depression and

anxiety in patients with type-2 diabetes. *Psychiatria Danubina*, 27(2), 179.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: van der Feltz-Cornelis C, Allen SF, Holt RIG, Roberts R, Nouwen A, Sartorius N. Treatment for comorbid depressive disorder or subthreshold depression in diabetes mellitus: Systematic review and meta-analysis. *Brain Behav*. 2021;11:e01981. <https://doi.org/10.1002/brb3.1981>